

Burn Dennet River and Tributaries Catchment Status Report 2008



Loughs Agency of the
Foyle Carlingford and
Irish Lights Commission



Burn Dennet River and Tributaries Catchment Status Report 2008



Report Reference LA/CSR/01/09

Headquarters

22, Victoria Road
Londonderry
BT47 2AB
Northern Ireland

Tel: +44(0)28 71 342100

Fax: +44(0)28 71 342720

general@loughs-agency.org

www.loughs-agency.org

Regional Office

Dundalk Street
Carlingford
Co Louth
Republic of Ireland

Tel+353(0)42 938 3888

Fax+353(0)42 938 3888

[carlingford@loughs-](mailto:carlingford@loughs-agency.org)
[agency.org](mailto:carlingford@loughs-agency.org)

www.loughs-agency.org

Inside cover picture of cock salmon in breeding
dress courtesy of Atlantic Salmon Trust



Written and Prepared by:
Art Niven
Rachel Scott
Declan Lawlor

Table of Contents

1.0	INTRODUCTION.....	6
	Fig 1.0 Main salmon producing rivers and tributaries in the Loughs Agency, Foyle and Carlingford areas.....	7
1.1	The Burn Dennet Catchment	8
	Fig 1.1 Burn Dennet River.	9
	Fig 1.11 Foyle and Carlingford catchments illustrating the main rivers of the systems.....	10
1.2	Atlantic Salmon and Sea Trout.....	11
1.3	Non Salmonid Fish Species.....	11
	Fig 1.3 Sample of fish from the Foyle estuary.....	11
2.0	ATLANTIC SALMON STOCKS	12
2.1	Redd Counts	12
	Table 2.1 Redd counts for Foyle system and Burn Dennet catchment 2004/05 – 2008/09.	12
	Fig 2.1 Redd counts for Foyle system and Burn Dennet catchment 2004/05 – 2008/09.....	13
	Fig 2.11 Redd counts for Foyle system 1952/53 – 2008/09	13
2.2	Juvenile Abundance	14
	Table 2.2 Loughs Agency semi-quantitative electrofishing classification system	15
	Fig 2.2 Burn Dennet River catchment salmon fry index 1998-2008..	15
	Fig 2.21 The mean abundance of salmon fry in 17 catchments in 2008	16
	Fig 2.22 Salmon 0+ electrofishing site classification 2008.....	17
3.0	TROUT STOCKS	18
	Fig 3.01 Burn Dennet River catchment trout fry index 1998-2008.....	18
	Fig 3.02 The mean abundance of salmon fry in 17 catchments in 2008	19
	Fig 3.03 Trout 0+ electrofishing site classification 2008.....	20
4.0	MARINE SURVIVAL	21
	Table 4.0 Marine survival rates for the River Bush of 1SW grilse 2002-2007	21
	Fig 4.0 Marine survey areas for salmon in 2008.....	22
	Fig 4a RV Celtic Explorer SALSEA research cruise 2008	23
	Figure 4b Picture from the Irish Research Vessel Celtic Explorer taken during the second SALSEA research cruise	23
	Figure 4c Rotary screw trap in position on the River Faughan	24
	Table 4.01 Numbers and average weight and length of salmon smolts tagged.....	24

Table 4.02 Recapture data from River Faughan CWT programme.....	25
Figure 4d Salmon smolt run timing and abundance from rotary screw trap sub sample	25
Table 4.03 Age class of salmon smolts migrating from the Faughan catchment in 2004	26
Fig 4.01. Atlantic salmon smolts, brook lamprey, river lamprey and sea lamprey.....	26
5.0 ADULT ABUNDANCE.....	27
5.1 Recreational Fisheries	27
Table 5.1 Declared rod catch returns for salmon and trout in the Foyle and Carlingford areas..	27
Table 5.11 Declared catch from the Burn Dennet catchment for salmon and sea trout 2001-2008	28
Fig 5.1 Recreational Fisher	28
5.2 Commercial Fisheries.....	28
Table 5.2 Declared catch from the commercial salmon fisheries 1998-2008	29
Fig 5.2 Commercial Fishing	29
5.3 Counters.....	29
Fig 5.3 Fish counting facilities at Sion Mills, River Mourne.....	30
Table 5.3 Sion Mills fish counter figures 2002-2008.....	30
Fig 5.31 Monthly fish counts on the River Mourne 2004-2008 as recorded at Sion Mills	30
Fig 5.32 Annual fish counts on the River Mourne 1987-2008	31
5.4 Conservation Limits/Spawning targets	31
Table 5.4 Upstream of Sion Mills estimated egg deposition 2002-2007	32
6.0 HABITAT MONITORING	32
Fig 6.0 Life cycle unit depicting the type of habitat found in spawning, nursery and holding	33
Fig 6.01 Examples of spawning, nursery and holding habitat	33
Fig 6.02 Burn Dennet catchment habitat survey	34
7.0 LAND USE	35
Fig 7.0 Burn Dennet catchment land use classification	35
Fig 7.01 Burn Dennet catchment land use classification map	36
8.0 WATER QUALITY.....	37
Fig 8.0 Loughs Agency chemical water quality testing in the laboratory	37
Fig 8.01 Burn Dennet catchment average suspended solids results 2008	40
Fig 8.02 Burn Dennet catchment Ammonia results 2008.....	41
Fig 8.03 Burn Dennet catchment phosphorous results 2008	42
Fig 8.04 Burn Dennet catchment Biological Oxygen Demand (BOD) results 2008.....	43
Fig 8.05 Burn Dennet catchment Biological Monitoring Working Party results 2008.....	44

Table 8.0 chemical GQA class limiting criteria	45
Table 8.01 Biological GQA class limiting criteria	46
Figure 8.06 Flattened mayfly nymph from the order <i>ephemeroptera</i>	46
8.1 Favourable Condition Tables, Target Levels	47
Table 8.1 Favourable condition targets for Atlantic salmon.....	47
Fig 8.1. Chemical General Quality Assessment (GQA) Burn Dennet catchment 2007	49
Fig 8.11 Biological General Quality Assessment (GQA) Burn Dennet catchment 2007	50
9.0 CONSERVATION AND PROTECTION	51
Table 9.0 Conservation and protection duties in the Burn Dennet catchment 2002-2008	51
Table 9.01 Seized nets, salmon, rod/reels and vehicles in the Foyle system 2002-2008.....	51
Table 9.02 Seizures in the Burn Dennet catchment 2006-2008	51
9.1 Habitat Improvement Works	52
Fig 9.1. Examples of in-stream habitat improvements on the Camus Burn	54
10.0 ENVIRONMENTAL ISSUES.....	54
11.0 DESIGNATED AREAS	55
12.0 GENETIC STUDY.....	56
13.0 POLLUTION MONITORING	56
14.0 FISHERIES OFFICERS BURN DENNET AREA REPORT 2008	57
14.1 Fishery Officers Report	57
15.0 ACTIONS FOR 2009.....	58
15.1 Foyle and Carlingford Areas Ongoing Actions for 2009	58
15.2 Burn Dennet Catchment Potential Habitat Improvement Schemes for 2009	59
15.3 Burn Dennet Catchment Specific Actions for 2009.....	60

Burn Dennet River and Tributaries Catchment Status Report 2008

1.0 INTRODUCTION

The annual Burn Dennet River and Tributaries Catchment Status Report has been updated to include results collated on the fishery resources within the catchment in 2008 and reports and discusses additional environmental information which when combined can facilitate in the sustainable management of aquatic resources within the catchment.

This is one of an annual series of catchment status reports produced by the Loughs Agency. The primary objective of the catchment status reports is to disseminate catchment specific information to all interested stakeholders. Over recent years the catchment status reports have been consulted widely by a variety of stakeholders including local angling associations, fishery owners, statutory bodies, environmental consultants and private individuals. The catchment status reports provide summary data which demonstrates the work that the Loughs Agency conducts within specific catchments and outlines catchment specific objectives.

The theme for the 2008 catchment status reports is aquatic and riparian habitats. Both habitats are essential for a balanced ecosystem ensuring that the biodiversity of wild salmonids and other native flora and fauna is preserved.

Significant focus has been placed on aquatic resource management by a combination of national, European and international legislation and conventions. In 2009 the first River Basin Management plans produced under the European Union Water Framework Directive will outline a structure for classifying all water bodies as having good ecological status. This will be an iterative process over many reporting cycles with significant improvements in water quality expected over coming years. Partnership development and collaborative working has been highlighted as an essential mechanism to achieve this and the Loughs Agency is playing an active role as a significant stakeholder and manager of aquatic resources within the Foyle and Carlingford areas.

In addition the European Union Habitats Directive and Birds Directive have resulted in the designation of Special Areas of Conservation (SAC) and Special Protection Areas (SPA). Together these designations have formed the NATURA 2000 network of protected sites. Throughout the Foyle and Carlingford areas there are significant areas designated as SAC or SPA. Under the SAC designation Atlantic salmon is one of the main species that led to the significant number of Foyle area tributaries being

designated. Designation provides an additional mechanism for the protection and conservation of these important aquatic ecosystems.

Northern Ireland and the Republic of Ireland are also a signatory through their membership of the European Union to the North Atlantic Salmon Conservation Organisation (NASCO) convention for the conservation of salmon in the North Atlantic Ocean. Contracting parties are required to contribute towards the compilation and implementation of focus area reports the most recent of which was on the Protection Restoration and Enhancement of Salmon Habitat. The Loughs Agency provided input for areas within its jurisdiction.




Figure 1.0 Main salmon producing rivers and tributaries in the Loughs Agency, Foyle and Carlingford areas. Foyle 1 Burn Dennet, 2 Camowen, 3 Culdaff, 4 Deelee, 5 Derg, 6 Drumragh, 7 Fairywater, 8 Faughan, 9 Finn, 10 Muff, 11 Glenelly, 12 Glenmornan, 13 Mourne, 14 Owenkillew, 15 Owenreagh East, 16 Owenreagh South, 17 Roe, 18 Strule, 19 River Foyle (tidal). Carlingford Only the Whitewater (10) has a consistent population of Atlantic salmon present. 1 Bessbrook, 2 Cassy Water, 3 Clanrye, 4 Ghann, 5 Greenore, 6 Killbroney, 7 Moygannon, 8 Newry, 9 Ryland, 10 Whitewater

Under the Foyle Fisheries Acts 1952 and the recent amendment by the Foyle and Carlingford Fisheries Act/Order 2007 there are various legislative requirements and offences including under the 2007 legislation a new offence of removal of any bed material from the freshwater portion of any river in the Foyle or Carlingford areas without consent of the Loughs Agency. This has provided an important control on activities likely to impact directly on aquatic habitats and indirectly on fish populations.

1.1 The Burn Dennet Catchment

The Burn Dennet River has many fast flowing streams which have eroded deep winding valleys carving the landscape into rounded, dome shaped hills. The landscape changes throughout the catchment and is predominantly composed of rolling farmland, steep wooded valleys and moorland capping the summits of higher hills. Glaciofluvial deposits have formed steeply undulating landforms on valley sides. The Burn Dennet River valley has a high aesthetic value which includes part of the Sperrin Mountains which forms the hilly backdrop to the Donemana basin. The valleys surrounding the Burn Dennet River and tributaries have pockets of semi-natural woodland.

Land use is dominated by improved pastures for grazing, rough grassland for sheep grazing and upland heathland. Woodland cover is low outside of the river valleys. Sand and Gravel pits are common on valley edges and in places have damaged the valley landscape. The Burn Dennet River and tributaries have a channel length of approximately 42 km and have a catchment area of 149km².

The Burn Dennet catchment is impacted upon by a wide range of anthropogenic factors within both the terrestrial and aquatic environments. A diverse array of impacts include amongst others; agriculture, sand and gravel extraction, commercial forestry, commercial and recreational fishing, industry, water abstraction, sewage treatment, diffuse and point source pollution, invasive plant species, urban sprawl and flood defences.

Increasing pressures on the aquatic environment within the Burn Dennet catchment requires appropriate monitoring, control and remediation if native biodiversity is to be preserved. The proximity of the Burn Dennet catchment to urban areas exacerbates many of these issues.

As the competent authority for fishery issues within the catchment the Loughs Agency are required to fulfil a variety of national and international obligations. European Directives including the Habitats Directive and Water Framework Directive and the transposing national legislation have assisted in creating a legislative framework in which to drive forward sustainable management of riparian and aquatic habitats and the native species which inhabit them.






Fig 1.1 Burn Dennet River.




Fig 1.11 Foyle and Carlingford catchments illustrating the main rivers of the systems and highlighting the Burn Dennet River and Tributaries

1.2 Atlantic Salmon and Sea Trout

Salmon and Sea Trout are referred to as being anadromous meaning that they migrate between the freshwater and marine environments returning to freshwater to reproduce. This complex life history exposes them to varied environmental pressures and recreational and commercial fisheries.

Adult Atlantic salmon return to their natal rivers where spawning takes place. Sea trout also demonstrate an ability to return to their natal river but their homing instinct may not be as strong as those of the Atlantic salmon. After the eggs hatch the juveniles (initially referred to as fry and then parr) remain in freshwater for up to three years.

Smoltification is the physiological adaptation which occurs when the juvenile salmon change from the parr stage (freshwater phase) to the smolt stage (marine phase). In the Foyle system this can occur after one, two or three years. Most Foyle salmon (referred to as post smolts) will remain after smoltification in the North Atlantic for one year and are referred to on their return to the coast and rivers as grilse. Salmon which stay at sea for longer than one year are referred to as multi sea winter (MSW) salmon.

1.3 Non Salmonid Fish Species

As highlighted earlier populations of other non salmonid fish species occur within the Burn Dennet catchment. In the past monitoring was targeted at salmonid species however with obligations under the Water Framework Directive other non salmonid fish species are being monitored more closely.

Fish species presence and abundance can act as a good environmental/ecological indicator demonstrating the ability of the aquatic habitat to support a diverse array of native species. Populations of the European Eel, Three spined stickleback, River/Brook and Sea Lamprey form an important part of the native fisheries biodiversity of the Burn Dennet catchment. Maintaining high standards of water quality and appropriate habitat for these species is essential for the overall health of the aquatic ecosystem.



Fig 1.3 Sample of fish from the Foyle estuary

2.0 ATLANTIC SALMON STOCKS

In order to describe the status of salmon stocks each of the following points need to be considered:

- Redd Counts
- Juvenile abundance
- Marine survival
- Adult abundance
- Exploitation

2.1 Redd Counts

Redds are spawning nests created by salmon or trout. Differentiation between salmon and trout redds can be made as salmon redds tend to be larger in size and trout tend to spawn earlier than salmon within the Foyle system. Research within the Foyle system using extensive annual redd count data has highlighted a good relationship between the number of redds and the total annual catch of salmon. Table 2.1 shows redd count data for the Burn Dennet catchment and the Foyle system with. Water flow is of significance when monitoring redds as in high water conditions the ability to see and count redds in rivers is impaired. Figure 2.1 outlines redd counts within the Foyle area and the Burn Dennet catchment.

Year	2004/05	2005/06	2006/07	2007/08	2008/09
Foyle System	2478	5354	1338	3039	5507
Burn Dennet Catchment	11	174	36	N/A	115
Burn Dennet as a % of Foyle	0.5	3	3	N/A	2

Table 2.1 Redd counts for Foyle system and Burn Dennet catchment 2004/05 – 2008/09.




Fig 2.1 Redd counts for Foyle system and Burn Dennet catchment 2004/05 – 2008/09. *Note 2006/07 had extremely poor water conditions for redd counting




Fig 2.11 Redd counts for Foyle system 1952/53 – 2008/09

2.2 Juvenile Abundance

Within the Loughs Agency jurisdiction trends in abundance of juvenile salmonids are monitored by annual semi-quantitative electrofishing surveys. The numbers, age and species of fish captured during five minute timed electrofishing surveys are compared with previous year's data allowing for change to be monitored, facilitating suitable fishery management practices to be implemented.

In 2008 a total of 403 sites were semi-quantitatively electrofished within the Foyle system. The results for each site for salmon and trout are classified as excellent (>25 fish), good (15-24 fish), fair (5-15 fish), poor (1-4 fish) and absent (0 fish), Table 2.2. Figures 2.2 - 2.22 outline the salmon 0+ electrofishing results and site classifications for the Burn Dennet catchment in 2008. Please consult previous status reports for site classifications in other years.

Semi-quantitative electrofishing was developed to monitor 0+ salmonids (fry/young of the year). In order to quantify the abundance of 1+ salmonids (parr and older) fully quantitative electrofishing surveys are required which can be used to calculate fish densities within a defined area. Rivers and tributaries with good environmental quality are more likely to support good populations of each year class.

Fish populations can vary considerably over time and location, it is therefore necessary to monitor the populations over a period of years to highlight meaningful trends before considering remedial activities such as habitat improvement works. These trends are being continually monitored by the Loughs Agency and the most appropriate management options considered.

There are a variety of reasons why electrofishing sites may be perceived to be under producing, these can include, lack of suitable juvenile habitat, the presence of impassable obstacles to migratory fish species on lower sections of a tributary, pollution, inconsiderate channel maintenance, tunnelling by bank side vegetation, stream gradient and poor forestry practices etc. The critical point is to recognise the major factors at play and to investigate all possible reasons for underproduction accepting that there may be inherent reasons as to why production may not be improved upon in certain areas. When the same areas are surveyed for other non salmonid species it may be discovered that they provide habitat more suited to these species. Habitat improvement works and the rationale behind them are discussed in greater detail later.

Obligations under the Water Framework Directive are driving quantitative surveys of both salmonid and non salmonid species under proposed Surveillance, Operational, Investigative and Protected Area monitoring programmes.






Symbol	Grade	Number of 0+ Salmonids
	Excellent	>25
	Good	15-24
	Fair	5-14
	Poor	1-4
	Absent	0

Table 2.2 Loughs Agency semi-quantitative electrofishing classification system for 0+ salmon and trout




Fig 2.2 Burn Dennet River catchment salmon fry index 1998-2008. Note number of sites surveyed has fluctuated annually, the number of sites surveyed has stabilised since 2002.




Fig 2.21. The mean abundance of salmon fry in 17 catchments in 2008 from semi quantitative electrofishing.




Fig 2.22 Salmon 0+ electrofishing site classification 2008

3.0 TROUT STOCKS

Annual trends in the populations of juvenile trout are also monitored within the Loughs Agency jurisdiction using the same methodology and classification system as those employed for salmon. The semi quantitative electrofishing results for trout fry in the Burn Dennet catchment and site classifications are displayed in Figs 3.0, 3.01, 3.02 & 3.03.



Fig 3.0 Electrofishing survey and trout parr




Fig 3.01 Burn Dennet River catchment trout fry index 1998-2008. Note number of sites surveyed has fluctuated annually, the number of sites surveyed has stabilised since 2002.




Fig 3.02. The mean abundance of salmon fry in 17 catchments in 2008 from semi quantitative electrofishing. Note Sites selected were biased towards salmon habitat




Fig 3.03 Trout 0+ electrofishing site classification 2008

4.0 MARINE SURVIVAL

The numbers of salmon that survive to return to the freshwater environment are greatly influenced by conditions in the marine environment. Climate change leading to changes in sea surface temperatures, prey abundance, high seas fishing, marine pollution, sub lethal levels of pollution and predation all have an effect on the Atlantic salmon and indeed other migratory fish species chances of survival.

Marine survival trends are monitored on a number of index rivers in the North East Atlantic where total trapping facilities are available for both migrating juvenile and adult populations. Total trapping allows for an accurate count of all migrant smolts (total freshwater production) and returning adults to be made and therefore an accurate estimate of marine survival. These projects are facilitated by the use of Coded Wire Tags (CWT). Coded wire tags are small (2-3mm long) micro tags that are injected automatically by a CWT device into the snout cartilage of anaesthetised fish remaining there for the duration of the life of the fish. CWT fish also have their adipose fin (small fin between the dorsal fin and caudal fin (tail fin)) removed so that they can be identified in the various fisheries that may intercept them. In Ireland a comprehensive screening programme is conducted at all major landing ports and markets. This programme is important in monitoring the effect of the remaining salmon fisheries on salmon stocks from rivers both within and outside of the island of Ireland.

Trends in marine survival for the River Bush (nearest index river to the Foyle system) confirm patterns observed elsewhere on the southern stocks of North Eastern Atlantic salmon, which indicate that marine survival can be variable between stocks and years. In the River Bush marine survival has decreased considerably over recent years as outlined in Table 4.0.

Year of Smolt Cohort	Year of Returning 1SW Grilse	Marine Survival %
Pre 1996	Pre 1998	Circa 30%
2002	2003	5.9
2003	2004	4.3
2004	2005	4.6
2005	2006	4.2
2006	2007	13.0
2007	2008	7.5

Table 4.0 Marine survival rates for the River Bush of 1SW grilse (after exploitation at sea) pre 1996 and 2002-2007 smolt cohort. Data supplied by Agri Food and Bioscience Institute, River Bush Salmon Research Station

The figures outlined in table 4.0 are mirrored by those for other index rivers monitoring the southern stocks of North Eastern Atlantic salmon populations. These figures suggest that salmon are facing increased pressure for survival at sea. A major new international research project called SALSEA - Merge has been developed by scientists from the North Atlantic Salmon Conservation Organisation (NASCO) parties and its research wing the International Atlantic Salmon Research Board (IASRB). There are twenty consortium members in total including the Loughs Agency. SALSEA aims to monitor how Atlantic salmon use the ocean; where they go; how they use ocean currents, and the ocean's food resources, and what factors influence migration and distribution at sea. Research cruises commenced in 2008 and will continue in 2009 to collect the necessary data to answer the questions listed above. Over 426 post smolts were caught by the two Irish cruises and 363 post smolts caught by the Faroese in the areas highlighted below. Further information and project details can be found at: <http://www.nasco.int/sas/salsea.htm>




Fig 4.0 Marine survey areas for salmon in 2008



Fig 4a RV Celtic Explorer SALSEA research cruise 2008



Figure 4b Picture from the Irish Research Vessel Celtic Explorer taken during the second SALSEA research cruise 16-24th May 2008

Since 2003 partial smolt trapping including CWT tagging has been conducted in the Faughan catchment using a rotary screw trap, Fig 4c.



Figure 4c Rotary screw trap in position on the River Faughan directly below the fish pass at Campsie barrage.

Smolt trapping can have a number of objectives including the monitoring of both salmonid and non salmonid species. Sampling of the age composition, obtaining information on run timing and recording length/weight data is conducted in tandem with the tagging programme. As mentioned above total counts of migrating smolts can be made on rivers. Where this is unfeasible due to the absence of total trapping facilities, total smolt migration can be estimated by means of a mark-recapture experiment.

In 2004 an estimate of total smolt production for the Faughan catchment was made by a mark-recapture study resulting in a minimum run size estimate of 33,854 migrating salmon smolts. The estimate was a minimum due to a number of high water events that prevented the smolt trap from fishing for a period of time during the peak smolt migration period. Tables 4.01 and 4.02 outline numbers of salmon smolts tagged from 2003-2008 and recapture data for 2003 and 2004.

Year	No of Salmon Smolts Tagged	Average Length (mm)	Average Weight (g)
2003	2113	149	33.45
2004	2500	134	24.6
2005	2210	133	23.6
2006	1025	133	25.36
2007	2062	135	27.1
2008	1865	130	22.1

Table 4.01 Numbers and average weight and length of salmon smolts tagged on the River Faughan 2003-2008. Coded Wire Tagging equipment was purchased by the Loughs Agency in 2005 with funding secured from the European Regional Development Fund through the INTERREG IIIA Programme, administered by the Environment and Heritage Service, on behalf of the Department of Environment.

Year Tagged	Year Recaptured	Numbers Recaptured	Recapture Location
2003	2004	12	Greencastle, Burtonport, Malin Head, Belmullet and Torr Head
2004	2005	16	Greencastle, Malin Head, Donegal and Galway Bay
2005	2006	3	Greencastle
2006	2007	2	Greencastle and Ballycastle

Table 4.02 Recapture data from River Faughan CWT programme. Data for fish tagged in 2007 and recovered in 2008 will not be available until 2009.




Figure 4d Salmon smolt run timing and abundance from rotary screw trap sub sample, River Faughan 2004-2008. Breaks in data are due to closure of trap during high water conditions

In 2004 a detailed examination was carried out on the age class of migrating salmon smolts in the Faughan catchment, Table 4.03.

Age at Smolting	%
1	13
2	83
3	4

Table 4.03 Age class of salmon smolts migrating from the Faughan catchment in 2004



Fig 4.01. From top to bottom, Atlantic salmon smolts from the Faughan Catchment, brook lamprey, river lamprey and sea lamprey also caught in the River Faughan smolt trap

5.0 ADULT ABUNDANCE

Adult Atlantic salmon abundance is assessed in three ways: directly by using commercial netting/recreational rod catches and fish counters and indirectly by reference to conservation limits/spawning targets.

Using catch data as a measure of population status is a well established and extensively used technique. In the Foyle system annual commercial and recreational catch data has been recorded since the establishment of the Foyle Fisheries Commission in 1952, with some data available before this period. The relationship between catch and stock is complex and care should be applied in interpretation. A more precise measure of catch incorporates fishing effort (number of licences issued or the amount of time fished) and is referred to as catch per unit effort (CPUE).

5.1 Recreational Fisheries

One problem encountered when analysing catch data is unreported catch. All recreational fishers are required by law to make catch returns. This information facilitates management decision making and therefore it is vitally important that all catch returns are accurate and made promptly at the seasons end.

Year	Declared Rod Catch Salmon	Declared Rod Catch Sea Trout	Returns as a % of Licences Issued
1999	1022	679	3.74
2000	723	417	2.55
2001	3188	450	17.68
2002	5117	1010	27.93
2003	1844	361	15.5
2004	2285	75	13.99
2005	4084	413	25.77
2006	3476	469	37
2007	4929	379	22.11
2008	4060	815	54.94

Table 5.1 Declared rod catch returns for salmon and trout in the Foyle and Carlingford areas. Note figures include the Clanrye and Whitewater in the Carlingford area from 2001 onwards. Carcass tagging was introduced in 2001.

Year	Declared Catch Burn Dennet Catchment Salmon	Declared Catch Burn Dennet Catchment Sea Trout
2002	7	48
2003	2	18
2004	6	10
2005	6	13
2006	1	0
2007	3	0
2008	18	32

Table 5.11 Declared catch from the Burn Dennet catchment for salmon and sea trout 2001-2008



Fig 5.1 Recreational Fisher

5.2 Commercial Fisheries

Commercial fisheries have traditionally operated within the Foyle sea area, Lough Foyle and tidal River Foyle. The drift net and draft net fisheries as well as the rod fisheries have been closely regulated with a real time management regime in place to monitor the numbers of fish migrating up key rivers. If predetermined numbers of fish have not been counted by the strategically placed electronic fish counters at Sion Mills weir (River Mourne), Campsie Barrage (River Faughan) and the Plumb Hole (River Roe) then specified closures of the commercial and/or recreational fisheries are enforced.

In 2007 new regulations were introduced to reduce the number of commercial nets operating within the Foyle area and all mixed stock interceptory drift nets seaward of Lough Foyle were curtailed. This decision was made to comply with the EU Habitats Directive, similar curtailment of mixed stock fisheries were introduced in the Republic of Ireland. Within the Foyle area this was achieved through a voluntary hardship scheme. 18 out of 112 drift nets remain in Lough Foyle, those remaining have been reduced in size from 900m to 500m and 10 out of 50 draft nets remain. This represents a significant reduction of netting effort. Regulations were also introduced to limit the numbers of fish which could be retained by the recreational rod fishery throughout the Foyle and Carlingford areas.

Year	Drift Catch	Draft Catch	Total Drift and Draft
1998	31296	11141	42437
1999	15397	7893	23290
2000	22333	10339	32672
2001	13500	9476	22976
2002	28851	11917	40768
2003	15741	16991	32732
2004	12800	9490	22290
2005	13391	12143	25534
2006	6160	6031	12191
*2007	2598	2774	5372
2008	1248	2924	4172

Table 5.2 Declared catch from the commercial salmon fisheries 1998-2008. Note 100% rate of catch returns. * Reduced numbers of commercial nets operating in the Foyle area from 2007



Fig 5.21 Commercial Fishing. Draft netting on the tidal River Foyle and drift netting in Lough Foyle

5.3 Counters

Within the Foyle system a number of river catchments have electronic fish counting facilities that provide estimates on the run timing and abundance of fish >45cm. The Burn Dennet catchment does not have a fish counter at present. In the absence of a counter the fisheries within the Burn Dennet catchment are controlled by monitoring the run timing and abundance of fish passing over the Sion Mills fish counting facilities on the River Mourne. A time series of counts for the Sion Mills fish counting station on the River Mourne is outlined in table 5.3. The Sion Mills fish counting station counts fish destined for all rivers located upstream of this point and is used to control both commercial and recreational fishing within the entire Foyle system.



Fig 5.3 Fish counting facilities at Sion Mills, River Mourne

Year	Number of fish >45cm
2002	12991
2003	12129
2004	10270
2005	9397
2006	9926
*2007	*3714
*2008	*3452

Table 5.3 Sion Mills fish counter figures 2002-2008. *Note low counts in 2007 and 2008 were influenced by high water levels during peak run timing during June and July. Sion Mills is a partial counter and does not cover the entire length of the weir. In high water levels as experienced in 2007 and 2008 significant numbers of fish can bypass the counting channels. Methods to reduce this are being investigated as part of the Fish Counter Programme Review, commissioned in 2008.




Fig 5.31 Monthly fish counts on the River Mourne 2004-2008 as recorded at Sion Mills




Fig 5.32 Annual fish counts on the River Mourne 1987-2008

5.4 Conservation Limits/Spawning targets

Another way to assess adult salmon stock status is to monitor run sizes on rivers and to compare them with predefined reference points called conservation limits. In the Foyle system the conservation limits define a level of spawning that optimises the sustainable catch by commercial and recreational fisheries. If exploitation rates increase above the sustainable catch levels the catch may increase in the short-term but the stock will eventually reduce. Conservation limits demarcate the spawning stock level at which recruitment would begin to decline significantly (NASCO). The real time management regime incorporating the setting of management targets and spawning targets implemented in the Foyle aims to manage the fisheries and spawning populations in a sustainable manner. The management and spawning targets are set for the various river catchments based on the amount and quality of nursery habitat present. River habitat surveys are carried out along each stretch of river and graded according to the type and quality. Egg deposition levels are set according to the quality grading of each section of nursery habitat.

There are four grades of nursery habitat, however for the purpose of setting egg deposition levels only grades 1-3 are utilised. Grade 1 denotes the best quality habitat. The egg deposition rate/carrying capacity is set as follows. Grade 1 = 10 eggs per m², grade 2 = 5 eggs per m² and grade 3 2.5 eggs per m². The total number of eggs is calculated by multiplying the area of each grade of nursery habitat by the

appropriate density of eggs per m². 25% is deducted from the management target allowing for loss of salmon by angling (15%) and poaching and predation (10%). The remaining figure is referred to as the conservation limit/spawning target.

Once the number of eggs required for each river has been established this can be converted to a total number of fish required to achieve the management targets and conservation limit/spawning targets. The average fecundity (number of eggs produced per female) of Foyle salmon has been estimated at 2500 and the ratio of female to male salmon estimated at 60:40. When combined with the amount of nursery habitat of the various grades this equates to the conservation limit/spawning target. A management target of 8000 adult Atlantic salmon has been set for above Sion Mills, this equates to a conservation limit/spawning target of over 6000 salmon.

Year	No of Fish Across Counter	Estimated Egg Deposition
2002	12991	14,614,875
2003	12129	13,645,125
2004	10270	11,553,750
2005	9397	10,571,625
2006	9926	11,166,750
*2007	*3714	*4,178,250
*2008	*3452	*3,883,500

Table 5.4 Upstream of Sion Mills estimated egg deposition 2002-2007. *Note 2007 and 2008 figures are minimum estimates of escapement due to high water levels resulting in the bypassing of the Sion Mills counting channels.

6.0 HABITAT MONITORING

The Loughs Agency has carried out extensive habitat surveys on all the major rivers and tributaries within the Foyle and Carlingford catchments. Habitat surveys are carried out on foot. Although time consuming this is at present the best method for classifying the various grades of habitat. Habitat is classified into one of three life cycle units Fig 6.0, the presence and order of which is essential to the productive capacity of a salmonid river. Other non salmonid species also benefit from diverse in-channel habitat. The life cycle unit categories include spawning, nursery and holding habitat. Each category is then graded on a scale of 1-4, 1 representing the best quality attainable and 4 the worst. Other data collected during these surveys include channel width and impassable barriers to migratory fish species.




Fig 6.0 Life cycle unit depicting the type of habitat found in spawning, nursery and holding zones



Fig 6.01 Examples of spawning, nursery and holding habitat




Fig 6.02 Burn Dennet catchment habitat survey

7.0 LAND USE

Land use classification is an important tool when assessing the potential impacts within a particular river catchment or indeed when looking at specific land use and land management practices. Land use impacts could have either a positive or negative impact on rivers and tributaries. A good understanding of the land use within a catchment is therefore imperative in managing at a catchment scale.

Land use in Northern Ireland has been captured using satellite imaging technology and classified to type. The following figures outline the broad land use classification within the Burn Dennet Catchment.




Fig 7.0 Burn Dennet catchment land use classification




Fig 7.01 Burn Dennet catchment land use classification map

8.0 WATER QUALITY

Routine water quality monitoring within the Foyle and Carlingford areas is conducted by the Northern Ireland Environment Agency (NIEA) of the Department of the Environment for Northern Ireland and the County Councils in the Republic of Ireland (Donegal County Council and Louth County Council). Routine sampling is conducted regularly for both chemical and biological General Quality Assessments (GQA).

In addition to the routine river monitoring carried out by the NIEA and the County Councils the Loughs Agency conducts proactive and reactive pollution investigations to investigate or highlight problems or potential problems which may have an effect on the aquatic environment and ultimately on the fish species and aquatic habitats.

In 2007 the Loughs Agency instigated a programme of monitoring at the tributary level for assessments of chemical and biological water quality. Four stations on tributaries of the Burn Dennet River were monitored for chemical water quality parameters including Biological Oxygen Demand (BOD), Suspended Solids, Ammonia and Phosphorous. Biological water quality was assessed using the Biological Monitoring Working Party (BMWP) a biotic scoring index.



Fig 8.0 Loughs Agency chemical water quality testing in the laboratory

The Loughs Agency also maintains a mobile pollution response unit containing aerating equipment and absorbent and non absorbent booms for oil and chemical spills. The unit can be rapidly deployed to the site of a pollution incident.

Water Quality Parameters

The following water quality parameters are monitored through the Loughs Agency monitoring programme and determined from water samples in the laboratory:

- Biochemical Oxygen Demand (BOD)
- Ammonia
- Phosphorus
- Suspended Solids

BOD

Any organic matter discharged into a river provides an immediate source of food for bacteria. These bacteria will break down the organic matter eventually into simple compounds such as carbon dioxide and water. Biochemical Oxygen Demand or Biological Oxygen Demand (BOD) is a chemical procedure for determining how fast biological organisms use up oxygen in a body of water. It is considered as an indication of the quality of a watercourse

Ammonia (NH₃)

Ammonia is generally found in small amounts in rivers and streams. This is due to microbiological activity and the resultant reduction of compounds containing nitrogen. High levels of ammonia can occur as a result from sewage pollution and have detrimental impacts on fish species.

Phosphorus (PO₄)

The over-loading of nutrients such as phosphorus in watercourses often leads to a process known as eutrophication. Eutrophication is a major environmental issue in Irish rivers and lakes. Sources of phosphorus include agricultural fertilizers and household detergents.

Suspended Solids

Particulate matter may be organic or inorganic in nature. Organic solids may consist of algal growths, indicative of eutrophic conditions. Inorganic solids generally are the result of discharge washings from sand and gravel extraction activities or quarries. Suspended solids can affect plant growth and fish habitats.

The following parameters are also recorded at each sample station by means of an electronic measuring probe:

- pH
- Temperature
- Dissolved Oxygen
- Conductivity

pH

This is a measure of the hydrogen ion concentration of a solution and therefore an indication of whether a liquid is acid or alkaline. The pH scale ranges from 0 (very acid) to 14 (very alkaline), with results generally influenced by geological conditions. Fish can be susceptible to changes in pH. Low pH levels are generally found in catchments with high forestry operation impacts.

Temperature

The effect of changes in temperature on living organisms, such as fish, can be critical. Thermal discharges from urban and industrial sources can lead to temperature increases in watercourses and increased stress on aquatic habitats and associated species.

Dissolved Oxygen

Sufficient levels of oxygen saturation in fresh waters are generally an indication of good ecological status and ideal for fish life. The main point to remember about oxygen solubility is that it has an inverse relationship with temperature. This helps explain why DO levels are generally lowest during summer low flow conditions, increasing the risk of pollution from discharges at this time.

Conductivity

The conductivity or electrical conductivity of a watercourse is a measure of its ability to conduct an electric current. Electrical conductivity estimates the amount of total dissolved salts, or the total amount of dissolved ions in the water. Electrical Conductivity is controlled by geology and any variations may be sourced to increased ions from wastewater from sewage treatment plants or urban run-off from roads.




Fig 8.01 Burn Dennet catchment average suspended solids results 2008. Values are in mg/l




Fig 8.02 Burn Dennet catchment Ammonia results 2008. Values are in mg/l




Fig 8.03 Burn Dennet catchment phosphorous results 2008. Values are in mg/l




Fig 8.04 Burn Dennet catchment Biological Oxygen Demand (BOD) results 2008. Values are in mg/l




Fig 8.05 Burn Dennet catchment Biological Monitoring Working Party results 2008

NIEA routinely monitor both the chemical and biological water quality within the rivers of Northern Ireland. In relation to chemical monitoring an extensive network of sampling stations are monitored for a variety of chemicals. The General Quality Assessment (GQA) is defined by limits for the concentrations of Biological Oxygen Demand (BOD), ammonia and dissolved oxygen (DO). The measures listed are indicators of the affect on water quality by waste water discharges and agricultural run-off containing organic material. Water quality can be affected by a variety of sources and the GQA determinands provide a recognised assessment of water quality.

The overall GQA class assigned to a section of river is based on the worst performing of the three measures (BOD, ammonia and DO). Table 8.0 outlines the standards for the chemical GQA.

GQA Class	Dissolved Oxygen (% Sat) 10- percentile	BOD (mg/l) 90-percentile	Ammonia (mg/l) 90-percentile
A (Very Good)	80	2.5	0.25
B (Good)	70	4	0.6
C (Fairly Good)	60	6	1.3
D (Fair)	50	8	2.5
E (Poor)	20	15	9.0
F (Bad)	<20	-	-

Table 8.0 chemical GQA class limiting criteria

The above table can be summarised as follows: for BOD and ammonia the section of river should contain less than the stated levels for at least 90% of the time. DO levels must not fall below the stated levels for more than 10 percent of the time.

In relation to biological monitoring an extensive network of sampling stations is also routinely monitored. The biological GQA is defined by observed measures of the abundance and diversity of macro invertebrates (for example freshwater shrimps, insect larvae and molluscs) compared to expected values as derived from a UK computer model adapted for Northern Ireland called River Invertebrate Prediction and Classification System (RIVPACS)

Different species of macro invertebrates are more sensitive to specific forms of pollution and therefore environmental quality indices (EQIs) based on biological results may be used to assess water quality. Macro invertebrates are also the dominant prey of both salmonid and some non salmonid fish species. The measure of diversity of a macro invertebrate community can be a more reliable indicator of the pollution pressures within a catchment than relying solely on an assessment of chemical water quality. The impacts of pollution on a macro invertebrate community are longer lasting and can highlight intermittent pollution impacts that may be missed through chemical water quality monitoring.

Biotic scoring systems have been developed to assign a score based on a standardised system to each sample site. One such system is the Biological Monitoring Working Party (BMWP). Generally the higher the BMWP score the better quality of the macro invertebrate community which reflects better water quality.

Based on a combination of biotic scoring systems biological GQA classes are assigned to sections of river. The two EQIs used are as follows:

$$\text{EQI}_{\text{taxa}} = \frac{\text{BMWP Observed Number of Taxa}}{\text{BMWP Predicted Number of Taxa (as derived from UK model)}}$$

BMWP Predicted Number of Taxa (as derived from UK model)

$$\text{EQI}_{\text{ASPT}} = \frac{\text{BMWP Observed ASPT (Average Score Per Taxon)}}{\text{BMWP Predicted ASPT (as derived from UK model)}}$$

BMWP Predicted ASPT (as derived from UK model)

Biological Class	EQI for ASPT	EQI for Taxon
A (Very Good)	1.00 or above	0.85 or above
B (Good)	0.90-0.99	0.70-0.84
C (Fairly Good)	0.77-0.89	0.55-0.69
DC (Fair)	0.65-0.76	0.45-0.54
E (Poor)	0.50-0.64	0.30-0.44
F (Bad)	<0.50	<0.30

Table 8.01 Biological GQA class limiting criteria




Figure 8.06 Flattened mayfly nymph from the order *ephemeroptera* high scoring macro invertebrate indicative of good water quality

European Council Directive 92/43/EEC of the 21st of May 1992 on the Conservation of Natural Habitats and on Wild Flora and Fauna (Also known as the Habitats Directive) was enacted in Northern Ireland under the European Communities Nature Conservation (Natural Habitats etc.) Regulations (Northern Ireland) 1995.

This indicates that those areas designated as areas of nature conservation designated for salmon should strive to achieve the water quality targets that are necessary for the designated species, which has additions to the GQA standards.

While it is current government policy for all rivers to meet the General Quality Assessment Standards, the Agency feels that favourable conditions standards as detailed below should be the water quality targets for all salmonid rivers within its jurisdiction.

8.1 Favourable Condition Tables, Target Levels

Natural Heritage of Northern Ireland Environment Agency have suggested guidelines for the determination of water quality, the first being the proposed UK Guidance on Conservation Objectives from monitoring designated sites and includes the following, which are considered as the favourable conditions tables.

They recommend Biological GQA Class A or B with no drop in class from the existing station, and Chemical GQA Class A or B depending on which type. This is in addition to no drop in class from the existing station. In addition to these favourable conditions tables, based on publications from Conserving Natura 2000 Rivers, the European Life Series, Ecology Series; No 7 Ecology of the Atlantic Salmon, *Salmo Salar* L. these publications have indicated that there are specific favourable conditions for this species.

An annual mean of less than 10 milligrams per litre suspended solids for nursery grounds, and annual mean of less than 25 milligrams per litre for migratory passage and the setting of soluble reactive phosphorous targets in relation to river reach types which should be as near background levels.

Parameter	Level	Percentile	Reason
BOD mg/l	2.5	90	GQA class A
Ammonia mg/l	0.25	90	GQA class A
Dissolved Oxygen % Saturation	80	10	GQA class A
Unionised Ammonia mg/l	0.025	95	Favourable Conditions Habitat Forming
Suspended solids mg/l			Specific for Atlantic Salmon
Nursery grounds	10		
Migratory passage	25		
Soluble Reactive Phosphorous mg/l	Background	-	Specific for Atlantic Salmon

Table 8.1 Favourable condition targets for Atlantic salmon

The Water Quality data in the reports has come from the Northern Ireland Environment Agency, Water Management Units Water Quality Archive. It is accepted by the Agency that monitoring is designed to ensure that water quality is monitored to ensure compliance with European Union directives. The monitoring however does not tie in well with the habitat and electrofishing survey monitoring carried out by the Loughs Agency, and as such the Loughs Agency instigated its own monitoring programmes in 2007 to link fish life, macro invertebrates and water quality into one holistic site evaluation. Additionally Northern Ireland Environment Agency, Water Management Unit data is not released in real-time and the data displayed below is for 2007, where the Loughs Agency status report is for 2008. By collecting and analysing water quality data the Loughs Agency can react to local water quality issues more effectively.

From 2008 water quality information from NIEA will be reported in a different manner. The European Union Water Framework Directive requires that numerous water quality parameters in addition to those traditionally monitored should be combined to provide an overall classification for water bodies. Additional parameters will include fish, diatoms and morphology amongst others. The Loughs Agency will monitor fish for WFD purposes and provide this information to NIEA.

The Loughs Agency will report on WFD classifications in the 2009 catchment status reports.




Fig 8.1. Chemical General Quality Assessment (GQA) Burn Dennet catchment 2007. Data supplied by NIEA




Fig 8.11 Biological General Quality Assessment (GQA) Burn Dennet catchment 2007. Data supplied by NIEA

9.0 CONSERVATION AND PROTECTION

The Loughs Agency continues to carry out an active fishery protection role throughout the catchments of the Foyle and Carlingford areas including the sea area, River Foyle and on all tributaries. Tables 9.0 and 9.01 outline the number of patrols and some duties carried out by the Loughs Agency staff in the Burn Dennet catchment and seizures for the Foyle area.

A team of Fishery Officers are responsible for the Burn Dennet catchment splitting their time between the Roe catchment the Faughan catchment, Glenmornan catchment, Moor Lough, Lough Ash and Binevenagh. This is in addition to regular fishery protection patrols on the River Foyle.

Year	No of Patrols	No of Licence Checks	Joint Patrols	On-site Inspections
2002	10	19	0	63
2003	13	12	0	22
2004	12	19	0	19
2005	10	21	1	32
2006	17	11	0	22
2007	27	5	2	12
2008	31	36	0	16

Table 9.0 Breakdown of conservation and protection duties in the Burn Dennet catchment 2002-2008

Year	2008	2007	2006	2005	2004	2003	2002
Nets	114	100	97	114	181	198	207
Salmon	92	56	91	118	130	155	94
Rod & Reel	136	85	26	10	16	12	22
Vehicles	1	0	2	1	1	0	0

Table 9.01 Seized nets, salmon, rod/reels and vehicles in the Foyle system 2002-2008

Year	Nets	Salmon/Trout	Rod/Reel	Vehicles	Boats
2006	0	0	5	0	0
2007	0	0	13	0	0
2008	0	0	8	0	0

Table 9.02 Seizures in the Burn Dennet catchment 2006-2008

9.1 Habitat Improvement Works

In addition to the traditional protection duties carried out by the Loughs Agency staff conservation and improvement of habitat has been increasing over recent years.

Over time man has imposed significant changes on the natural courses of many rivers and flood plains. The driving forces behind these changes have included amongst others; arterial drainage schemes to provide more suitable land for agricultural purposes, urban sprawl, infrastructure expansion (roads etc.), flood defences, water abstraction and hydro power generation. All have had a significant impact on the natural meanderings and discharges of rivers and tributaries resulting in faster runoff of floodwaters ultimately leading to a change in the morphology and flow regime of rivers and resultant impacts on fisheries.

While all these processes have had some impact within the Foyle system, it is considered to be a relatively natural system with natural river structure present in the catchments headwaters. In areas that have been altered methods for reinstating lost habitat are investigated and where appropriate action taken.

In 2006 the Loughs Agency carried out habitat improvement works within the Burn Dennet Catchment on the Camus Burn. A series of habitat Units were created in-channel, riparian fencing at an appropriate distance from the water course was erected, cattle passage constructed over the Burn and spawning gravel introduced above vortex weirs. The pictures below outline some of these features. This site has subsequently featured on RTE Eco-Eye and UTV Lesser Spotted Ulster. The Loughs Agency see this project as a good example of partnership working to improve aquatic and riparian habitat and regularly lead guided visits to promote this exemplar project to groups interested in carrying out similar works within the Foyle and Carlingford areas.








Fig 9.1. Examples of in-stream habitat improvements on the Camus Burn

10.0 ENVIRONMENTAL ISSUES

Some environmental issues affecting water quality have already been outlined previously. The following list presents some of the main habitat pressures to salmonids within the Foyle system:

- Agricultural activities – enrichment from natural and artificial fertilisers often make their way into watercourses, enhancing problems with eutrophication.
- Forestry activities – planting and felling operations can lead to increased loading of suspended solids in watercourses. Established forestry as a major upland land use has been attributed to increased acidification.
- Barriers to migration – a range of natural and anthropogenic features on rivers can lead to barriers for migrating salmonids and other fish species. These can include weirs and hydro-electric schemes.
- Gravel removal – gravel is extremely important for the creation of redds for spawning fish. Removal of gravel from the river bed in sensitive areas can destroy potential spawning and nursery habitat.
- Quarrying activities – the extraction of aggregates such as rock, sand and gravel has the potential to cause increased levels of suspended solids in

nearby watercourses. Sufficient mitigation measures should be in place at such sites to trap increased sediment loads entering rivers and streams.

- Abstraction – water abstraction from watercourses for a range of uses is increasing throughout the Foyle and Carlingford catchments. Unless appropriately assessed and licenced, these activities have the potential to reduce residual flow levels and alter the ecological status of our rivers. This is even more concerning in the light of climate change.
- Peat harvesting – Peat harvesting still occurs in small upland pockets throughout the Foyle system. It has the potential to increase sediment loading in receiving waters.
- Sewage treatment – sewage and waste water treatment works are under considerable pressure with the increase in urban development in our towns and villages. Several inadequate systems throughout the Foyle system continue to pollute rivers.
- Hydropower – small-scale hydropower schemes are beginning to appear on rivers throughout the Foyle and Carlingford catchments. Baseline fishery data must be provided to allow for sufficient assessment of any proposed scheme, unless located above an impassable fish barrier.
- Urban development – the expansion of large-scale housing developments and the associated pressures on waste water and sewage treatment works are a potential source of water pollution in the event of overflows.
- Drainage and canalisation – these have direct impacts on the quality of available fishery habitat within the catchments. Canalisation in particular can lead to the removal of important spawning, nursery or holding areas of rivers.
- Industrial discharges – larger urban areas with industrial discharges have the potential to cause pollution through toxic discharges and can alter the temperature of the watercourse.
- Septic tanks – a proliferation of single dwellings and their septic tanks is an ongoing area of concern. Initial research from parts of the Foyle system indicate that this is major contributor to decreased water quality and local increases in suspended solids.

11.0 DESIGNATED AREAS

The European Commission Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (EU Habitats Directive 92/43/EEC) requires that all member states designate Special Areas of Conservation (SACs) in order to protect threatened habitats and species. The European Commission Directive on the Conservation of Wild Birds (Birds Directive 79/409/EEC) also requires the designation of Special Protected Areas (SPA's). Together the designated SAC's and SPA's create the NATURA 2000 network of protected sites. A number of rivers have been designated as SAC's both in Northern Ireland and in the Republic of Ireland

including the River Foyle and selected tributaries. The Burndennet River however does not form part of this designation.

While not a designated SAC the Burn Dennet catchment has a number of nationally designated areas including; Corbylin Wood and Silverbrook Wood which have been designated as Areas of Special Scientific Interest (ASSI). ASSIs are areas of land that have been identified as being of the highest degree of conservation value. Corbylin Wood and Silverbrook Wood have been designated for upland Oakwood. A section of the headwaters of the Burndennet catchment are within the Sperrin Area of Outstanding Natural Beauty (AONB).

12.0 GENETIC STUDY

A baseline genetic survey was carried out in the Foyle system in 2003 and a resurvey conducted between 2006 and 2008 to analyse the populations of Atlantic salmon present within the Foyle catchment. Results confirmed the existence of genetically distinct populations between and within the rivers and tributaries of the Foyle area. An understanding of these genetically differentiated populations is required to facilitate appropriate management of conservation measures and the commercial/recreational fisheries.

The report concluded that genetic diversity is high between and within the various salmon populations present in the Foyle system. Each population has evolved over time creating distinct populations (with some gene flow from straying fish) that are best suited to the conditions present in a particular river or tributary. The non-uniform nature of the populations adds to the diversity of life history strategies exercised by Foyle salmon. Distinct differences such as run-timing and age at smolting can act as nature's insurance policy to any catastrophic events which would threaten a homogenous population.

The report stated that the current genetic structure and diversity of Foyle salmon is representative of what might be regarded as the native structure of wild salmon populations. The maintenance of genetic diversity is a core requirement for the long-term sustainability of wild populations, preserving the biodiversity of the wild salmonids of the Foyle system is therefore a primary objective of the Loughs Agency.

13.0 POLLUTION MONITORING

The Loughs Agency has a statutory obligation to monitor the pollution of watercourses. In conjunction with the Northern Ireland Environment Agency all reported pollution incidents are investigated.

14.0 FISHERIES OFFICERS BURN DENNET AREA REPORT 2008

In 2008 Fishery Officers Emmett Carten, David Robinson and David O'Brien continued to work within the Burn Dennet catchment. Their report follows.

14.1 Fishery Officers Report

During 2008 the Eastern Crew patrolled the Burn Dennet catchment with a mix of specialised and routine patrols. Potential threats to the fishery were identified and clearly monitored.

A wet 2008 posed additional pressures on both environmental and angling fronts.

Substantially higher water levels did provide enhanced angling conditions with an increase of reported catch. New carcass tagging and bag limit regulations were well received and adhered to by the angling fraternity. Reports of illegal angling were very low but the Eastern Crew continued to investigate all reports even though reports were largely dated and often inaccurate. Any reports of illegal netting centered around the confluence with the River Foyle. Routine patrols by foot and boat were regularly carried out.

The wet weather encountered in 2008 had the potential to create increased threats to water quality within the Burn Dennet catchment. However reports of pollution were reduced from 2007 levels and the Eastern Crew carried out both reactive and proactive investigations with specific attention paid to the sand plants within the catchments. Again a close working relationship with the Northern Ireland Environment Agency water quality team was maintained throughout 2008. Routine water quality sampling was carried out and analysis of the samples completed by the Loughs Agency laboratory throughout the summer months.

With reference to in stream work the project carried out on the Camus Burn was found to have bedded in well and became somewhat of an enhancement showcase for the Loughs Agency. Low water conditions led to a full walk through of the Burn Dennet catchment at spawning time with increased spawning activity recorded over 2007 figures.

A wet summer in 2008 led to a protracted electrofishing field season however hard work by the Eastern Crew often during unsocial hours led to successful completion of all the designated electrofishing stations on the Burn Dennet with healthy populations recorded.

15.0 ACTIONS FOR 2009

In order to fully utilise the extensive data resources collected and held by the Loughs Agency on the fish populations and habitats of the Burn Dennet catchment it is necessary to focus attention on specific management objectives.

The Loughs Agency has stated in its corporate plan 2008-2010 that it will conserve, protect, manage and improve the fisheries of the Foyle and Carlingford areas. By way of fulfilling these objectives a targeted series of actions utilising data collected over recent years will be implemented. Fishery owners and local angling clubs will continue to be consulted regarding any proposed works and stakeholder input sought.

15.1 Foyle and Carlingford Areas Ongoing Actions for 2009

Good water quality is essential for the conservation of productive aquatic ecosystems. Fish populations rely on unpolluted water for survival and feeding. The Loughs Agency is committed to ensuring deleterious matter does not enter any watercourse. Routine monitoring is conducted throughout the Foyle and Carlingford areas. Proactive pollution visits and water quality monitoring will continue in 2009.

Water quantity is becoming an increasingly important issue from a fisheries management perspective with continuing demand from a variety of sources including industry, hydro power generation and abstraction for meeting the ever growing needs of industry and the wider population. The Loughs Agency are aware of the conflicting needs of aquatic environments and water resource users and comment on development issues which may have an impact on the important aquatic resources of the Foyle and Carlingford areas with reference to national and international obligations.

In-channel and riparian habitat improvement projects provide an important mechanism by which to improve and protect valuable fishery resources. Over recent years the Loughs Agency has developed a number of projects designed to improve the survival and production of robust populations of juvenile salmonid and other native fish species. These programmes will continue where funding is available, The Loughs Agency also encourages local stakeholder groups to source appropriate funding to develop collaborative habitat improvement projects. The Loughs Agency can provide advice and recommendations for in-channel and riparian improvements and are eager to facilitate the development of such programmes.

Work is continuing to assess and record all **Barriers to Migration** within the catchments of the Foyle and Carlingford areas and these will be incorporated into

the Loughs Agency Geographical Information System (GIS). Where finances are available the removal of artificial barriers will be investigated.

Predation by cormorants and seals of economically important fish species continues to be a contentious issue. The Loughs Agency will continue to promote the development of a management strategy incorporating economic, social and environmental factors.

The Loughs Agency will continue to monitor the salmon and inland fishery resources of the Foyle and Carlingford areas, utilising best practice methods including fish counters, juvenile population surveys and catch returns. The importance of the Atlantic salmon resource has been further highlighted by recent genetic studies which have identified the presence of genetically distinct populations of salmon between and within main river catchments. This information will be utilised when developing habitat improvement programmes to ensure the presence of a diverse resource capable of withstanding change.

Invasive species in both aquatic and riparian habitats have become an important issue in fisheries management and in wider environmental management. Invasive species have the potential to significantly alter ecosystems and their function. The Loughs Agency is contributing towards the development and implementation of invasive species codes of practice.

15.2 Burn Dennet Catchment Potential Habitat Improvement Schemes for 2009

- Burn Dennet – upstream of Tullyard Bridge – Area lacks suitable nursery habitat especially parr water – possible provision of rubble mats and boulders.
- Burn Dennet – Down stream of Presbyterian Bridge – Lacks suitable spawning – possible measures to be put in place to enhance spawning and to make use of gravel that is currently exposed mid channel as a shoal during low water conditions
- Burn Gibbagh – possible reassessment of drainage flap barrier to permit upstream migration
- Burn Dennet – upstream of Donemana Road Bridge to Mill Pottery area currently suffers from tree/bush/shrub tunnelling. Recommend a programme of sensitive tree/branch cutting to remove this overhang and to create better light penetration to the watercourse

15.3 Burn Dennet Catchment Specific Actions for 2009

- Maintain high standards of conservation and protection within the Burn Dennet catchment
- Target all areas/individuals brought to Loughs Agency attention
- Implement habitat improvement schemes as dictated by business plan/corporate plan
- Conduct annual fish population surveys and spawning surveys
- Conduct ongoing water quality monitoring and investigate areas highlighted as being of concern
- Develop potential habitat improvement projects including riparian buffer zone creation, fencing, native species planting and in-channel habitat improvements including spawning bed and nursery habitat improvement
- Monitor forestry operations adjacent to watercourses or areas likely to impact on watercourses
- Assist with Water Framework Directive fish monitoring programme
- Monitor all sand and gravel extraction sites and onsite water management practices
- Ensure all fish passes, dams and mill races meet required standards