

# Carlingford Tributaries Catchment Status Report 2008



Loughs Agency of the  
Foyle Carlingford and  
Irish Lights Commission



# Carlingford Tributaries Catchment Status Report 2008

---



Report Reference LA/CSR/C001/09

---

## Headquarters

22, Victoria Road  
Londonderry  
BT47 2AB  
Northern Ireland

Tel: +44(0)28 71 342100

Fax: +44(0)28 71 342720

[general@loughs-agency.org](mailto:general@loughs-agency.org)

[www.loughs-agency.org](http://www.loughs-agency.org)

## Regional Office

Dundalk Street  
Carlingford  
Co Louth  
Republic of Ireland

Tel+353(0)42 938 3888

Fax+353(0)42 938 3888

[carlingford@loughs-agency.org](mailto:carlingford@loughs-agency.org)

[www.loughs-agency.org](http://www.loughs-agency.org)

Inside cover picture of Atlantic salmon kelt  
courtesy of Atlantic Salmon Trust



Written and Prepared by:  
Art Niven  
Rachel Scott  
Declan Lawlor

---

# Table of Contents

---

1.0	INTRODUCTION.....	7
	Figure 1. Main salmon producing rivers and tributaries in the Loughs Agency, Foyle and Carlingford areas.....	8
1.1	The Carlingford Area and Tributaries.....	9
	Fig 1.1 Carlingford area looking east from the Newry River estuary.....	10
	Fig 1.11 Carlingford area looking west from the Newry River estuary towards Newry .....	11
	Fig 1.12 Foyle and Carlingford catchments illustrating some of the main tributaries. ....	11
	Fig 1.13 Carlingford area and tributaries. ....	12
1.2	Atlantic Salmon and Sea Trout.....	13
1.3	Non Salmonid Fish Species.....	13
	Fig 1.3 Bream specimens recovered during a coarse fish survey of Newry Canal in 2002.....	13
2.0	ATLANTIC SALMON STOCKS.....	14
2.1	Redd Counts .....	14
2.2	Juvenile Abundance .....	14
	Table 2.2 Loughs Agency semi-quantitative electrofishing classification system .....	15
	Fig 2.21 Electrofishing on the Clanrye River and salmon parr .....	15
	Fig 2.22 Carlingford Area fry index 2000-2008 .....	16
	Fig 2.23 White Water catchment fry index 2000-2008.....	17
	Fig 2.24 Clanrye River catchment fry index 2000-2008 .....	17
	Fig 2.25 Salmon 0+ electrofishing site classification 2008.....	18
3.0	TROUT STOCKS.....	19
	Fig 3 Electrofishing survey and trout parr.....	19
	Fig 3.1 Carlingford Area fry index 2000-2008. ....	19
	Fig 3.11 Whitewater trout fry index 2000-2008. ....	20
	Fig 3.12 Clanrye trout fry index 2000-2008. ....	20
	Fig 3.13 Trout 0+ electrofishing site classification 2007 .....	21
4.0	MARINE SURVIVAL .....	22
	Table 4 Marine survival rates for the River Bush of 1SW grilse.....	22
	Fig 4 Proposed marine survey areas for salmon in 2008.....	23
	Fig 4a RV Celtic Explorer SALSEA research cruise 2008 .....	24

Figure 4b Picture from the Irish Research Vessel Celtic Explorer .....	24
Figure 4c Rotary screw trap in position on the River Faughan .....	25
Table 4.1 Numbers and average weight and length of salmon smolts.....	25
Table 4.12 Recapture data from River Faughan CWT programme.....	26
Figure 4d Salmon smolt run timing and abundance from rotary screw trap sub sample .....	26
Table 4.13 Age class of salmon smolts migrating from the Faughan catchment in 2004 .....	27
Fig 4.14. From top to bottom, Atlantic salmon smolts from the Faughan Catchment, brook lamprey, river lamprey and sea lamprey .....	27
<b>5.0 ADULT ABUNDANCE.....</b>	<b>28</b>
5.1 Recreational Fisheries .....	28
Table 5 Declared rod catch returns for salmon and trout in the Foyle and Carlingford areas.....	28
Table 5.1 Declared catch from the Carlingford system for salmon and trout 2001-2008.....	29
Fig 5.11 Angler on the upper reaches of the Faughan River.....	29
5.2 Commercial Fisheries .....	29
Table 5.2 Declared catch from the commercial salmon fisheries 1998-2008 .....	30
Fig 5.21 Commercial Fishing .....	30
5.3 Counters.....	30
Fig 5.3 Fish Pass and counter construction on the Newry River.....	31
Table 5.3 Newry/Clanrye River fish counter figures. ....	31
Fig 5.32 Monthly fish counts on the Newry River.....	32
5.2 Conservation Limits/Spawning targets .....	32
<b>6.0 COARSE FISH STOCKS .....</b>	<b>33</b>
Figure DL 1. Temperature/DO profile in Derryleckagh Lough .....	34
Figure DL 2. Location of benthic gillnets (red circles), floating gillnet (white circle) and fyke nets (red squares) in Derryleckagh Lough .....	36
Table DL 1. Number of fish caught in each net in Derryleckagh Lough .....	36
Figure DL 3. Length frequency of pike in Derryleckagh Lough .....	37
Figure DL 4. Length frequency of eels in Derryleckagh Lough.....	37
Figure DL 5. Derryleckagh Lough survey 2008.....	38
Table 6 Fish survey results from Newry canal 2007 .....	39
Fig 6 Newry Canal fish stock assessment 2007.....	39
Table 6.1 Fish survey results from Newry canal 2001 .....	40
Table 6.2 Fish survey results from Newry canal 2002. ....	40
Fig 6.1 Left to right Perch, Pike and Bream.....	41



7.0	HABITAT MONITORING .....	41
	Fig 7 Life cycle unit.....	41
	Fig 7.1 Examples of spawning, nursery and holding habitat .....	41
8.0	LAND USE .....	42
	Fig 8 Carlingford Area land use classification .....	42
	Fig 8.1 Carlingford Area land use classification map .....	43
9.0	WATER QUALITY.....	44
	Fig 9.0 Loughs Agency chemical water quality testing in the laboratory .....	44
	Fig 9.1 Carlingford area average suspended solids results 2008.....	48
	Fig 9.2 Carlingford area Ammonia results 2008.....	49
	Fig 9.3 Carlingford area phosphorous results 2008.....	50
	Fig 9.4 Carlingford area Biological Oxygen Demand (BOD) results 2008. ....	51
	Fig 9.5 Carlingford area Biological Monitoring Working Party results 2008.....	52
	Table 9.1 chemical GQA class limiting criteria .....	53
	Table 9.2 Biological GQA class limiting criteria .....	54
	Figure 9.6 Flattened mayfly nymph .....	54
9.1	Favourable condition tables, target levels.....	55
	Table 9.1 Favourable condition targets for Atlantic salmon.....	55
	Fig 9.07 Chemical General Quality Assessment (GQA) Carlingford area 2007 .....	57
	Fig 9.08 Biological General Quality Assessment (GQA) Carlingford area 2007 .....	58
10.0	CONSERVATION AND PROTECTION .....	59
	Table 10. Breakdown of conservation and protection duties in the Carlingford area .....	59
	Table 10.1 Seized nets, salmon, rod/reels and vehicles in the Foyle and Carlingford areas.....	59
	Table 10.2 Seizures in the Carlingford area 2008 .....	59
10.1	Habitat Improvement Works .....	60
	Fig 10.4 Reinstatement of pool with some repair work to a weir on the Kilbroney .....	60
	Fig 10.5 Creation of spawning habitat on the Kilbroney .....	61
	Fig 10.6 Repair of weirs on the Kilbroney .....	61
	Fig 10.7 Repair of damaged weir on the Kilbroney.....	61
	Fig 10.8 Trimming of overhead cover and pool reinstatement on the Kilbroney .....	62
	Fig 10.8 Grading of bank on the Kilbroney .....	62
	Fig 10.9 Pool excavation and repair of weir on the Kilbroney.....	63
	Fig 10.10 Repair of weir on the Kilbroney.....	63
	Fig 10.11 Repair of weir and reinstatement of pool on the Kilbroney .....	63

11.0	ENIRONMENTAL ISSUES.....	64
12.0	DESIGNATED AREAS .....	65
13.0	GENETIC STUDY.....	65
14.0	POLUTION MONITORING .....	66
15.0	FISHERY OFFICERS CARLINGFORD AREA REPORT 2008 .....	66
16.0	WHITEWATER RIVER SALMONID POPULATION SURVEY 2008 .....	67
	Fig 16 Salmonid densities at Ballyardle Bridge 2008 .....	67
	Fig 16.1 Salmon fry and parr length weight relationships at Ballyardle bridge 2008 N = 89 .....	68
	Fig 16.2 Length Frequency distribution of Whitewater salmon fry and parr in 2008 N = 89 .....	68
17.0	ACTIONS FOR 2009.....	69
17.1	Foyle and Carlingford Areas Ongoing Actions for 2009 .....	69
17.2	Carlingford Area Specific Actions for 2009 .....	70

# Carlingford Tributaries Catchment Status Report 2008

---

## 1.0 INTRODUCTION

This is the third annual Carlingford Tributaries Catchment Status Report. It has been updated to include results collated on the fishery resources within the catchment in 2008 and reports and discusses additional environmental information which when combined can facilitate in the sustainable management of aquatic resources within the catchment.

This is one of an annual series of catchment status reports produced by the Loughs Agency. The primary objective of the catchment status reports is to disseminate catchment specific information to all interested stakeholders. Over recent years the catchment status reports have been consulted widely by a variety of stakeholders including local angling associations, fishery owners, statutory bodies, environmental consultants and private individuals. The catchment status reports provide summary data which demonstrates the work that the Loughs Agency conducts within specific catchments and outlines catchment specific objectives.

The theme for the 2008 catchment status reports is aquatic and riparian habitats. Both habitats are essential for a balanced ecosystem ensuring that the biodiversity of wild salmonids and other native flora and fauna is preserved.

Significant focus has been placed on aquatic resource management by a combination of national, European and international legislation and conventions. In 2009 the first River Basin Management plans produced under the European Union Water Framework Directive will outline a structure for classifying all water bodies as having good ecological status. This will be an iterative process over many reporting cycles with significant improvements in water quality expected over coming years. Partnership development and collaborative working has been highlighted as an essential mechanism to achieve this and the Loughs Agency is playing an active role as a significant stakeholder and manager of aquatic resources within the Foyle and Carlingford areas.

In addition the European Union Habitats Directive and Birds Directive have resulted in the designation of Special Areas of Conservation (SAC) and Special Protection Areas (SPA). Together these designations have formed the NATURA 2000 network of protected sites. Throughout the Foyle and Carlingford areas there are significant areas designated as SAC or SPA. Under the SAC designation Atlantic salmon is one of the main species that led to the significant number of Foyle area tributaries being

designated. Designation provides an additional mechanism for the protection and conservation of these important aquatic ecosystems.

Northern Ireland and the Republic of Ireland are also a signatory through their membership of the European Union to the North Atlantic Salmon Conservation Organisation (NASCO) convention for the conservation of salmon in the North Atlantic Ocean. Contracting parties are required to contribute towards the compilation and implementation of focus area reports the most recent of which was on the Protection Restoration and Enhancement of Salmon Habitat. The Loughs Agency provided input for areas within its jurisdiction.




Figure 1. Main salmon producing rivers and tributaries in the Loughs Agency, Foyle and Carlingford areas. Foyle 1 Burn Dennet, 2 Camowen, 3 Culdaff, 4 Deelee, 5 Derg, 6 Drumragh, 7 Fairywater, 8 Faughan, 9 Finn, 10 Muff, 11 Glenelly, 12 Glenmornan, 13 Mourne, 14 Owenkillew, 15 Owenreagh East, 16 Owenreagh South, 17 Roe, 18 Strule, 19 River Foyle (tidal). Carlingford Only the Whitewater (10) has a consistent population of Atlantic salmon present. 1 Bessbrook, 2 Cassy Water, 3 Clanrye, 4 Ghann, 5 Greenore, 6 Killbroney, 7 Moygannon, 8 Newry, 9 Ryland, 10 Whitewater

Under the Foyle Fisheries Acts 1952 and the Foyle and Carlingford Fisheries Act/Order 2007 there are various legislative requirements and offences including under the 2007 legislation a new offence of removal of any bed material from the freshwater portion of any river in the Foyle or Carlingford areas without consent of the Loughs Agency. This has provided an important control on activities likely to impact directly on aquatic habitats and indirectly on fish populations.

## 1.1 The Carlingford Area and Tributaries

Located in the North East of the island of Ireland and composed of significant areas of County Armagh, County Down and County Louth, the Carlingford catchments cover an area of approx 544 km<sup>2</sup>.

The catchments of the Carlingford area can be broadly divided into 4 main landscape types, the Newry basin, Slieve Roosley, Carlingford Lough and Mourne mountains. The Newry basin is characterised by a large scale rolling drumlin landscape (this drift geology is composed of deposits left by retreating glaciers) situated between the Ring of Gullion and the Mourne Mountains. The Newry basin is drained by the Newry/Clanrye River and tributaries. The main land use consists of improved pastures of good condition becoming increasingly rough on the fringes of the Mourne foothills. To the southeast drumlins are displaced by broader ridges separated by narrow, flat-bottomed valleys with ribbon loughs and bogs such as Derryleckagh Lake and Greenan Lough. The underlying solid geology is composed of basalt, sandstones and shales.

The Slieve Roosley landscape lies between Newry and the Mourne Mountains and is characterised by open, exposed hills with a rugged profile, which are dissected by a number of river valleys. The Rostrevor Glen and Killbroney River together form a marked feature along the eastern boundary of the area, which is underlain by a complex geology of igneous and sedimentary rocks. The hills are used for sheep grazing and are characterised by rough, open, unfenced pastures of moorland grasses, gorse, bracken and sedges. The fringes comprise semi-improved pastures of small fields enclosed by stonewalls and gorse hedgerows.

The Mourne Mountains Landscape (particularly associated with the Whitewater River catchment) is characterised by steep rock and scree covered mountain slopes capped with granite torrs, falling to the sea on their eastern edge. It is largely a wild upland landscape composed of exposed heath, thin grass cover, rock and scree slopes with rough grazing for sheep and some cattle. The underlying geology is dominated by granite which is reflected by the characteristic torrs which cap the mountain tops. Rocky mountain streams occupy the steep glens which dissect the mountain ridges.

Carlingford Lough is a low energy estuary filling a structurally controlled (NW-SE fault) glacially scoured depression. The estuary mouth is shallow which allows wave focusing of southwesterly storms onto the northern shoreline where erosion has left a number of bays dominated by gravel beaches.

Carlingford Lough supports a range of unusual and rich littoral/shoreline communities, including sheltered sands, muddy sands, muds and boulder shores. It exhibits a good natural transition from lower shore communities, through upper shore saltmarsh to fen vegetation. Mill Bay in particular supports the largest intact block of

saltmarsh in Northern Ireland and the area is internationally important in terms of numbers of wildfowl and waders that over-winter on the site.

The Carlingford area and tributaries are impacted upon by a wide range of anthropogenic influences within both the terrestrial and aquatic environments. A diverse array of impacts include amongst others; agriculture, aquaculture, sand and gravel extraction, quarrying, commercial forestry, commercial and recreational fishing, industry, water abstraction, sewage treatment, diffuse and point source pollution, invasive plant species, urban sprawl, flood defences and heavily modified water bodies. Increasing pressures on the aquatic environments within the Carlingford area and tributaries requires appropriate monitoring, control and remediation if native biodiversity is to be preserved and enhanced. The proximity of some of the Carlingford area and tributaries to a large urban area exacerbates many of these issues.

As the competent authority for fishery issues within the catchment the Loughs Agency are required to fulfil a variety of national and international obligations. European Directives including the Habitats Directive and Water Framework Directive and the transposing national legislation have assisted in creating a legislative framework in which to drive forward sustainable management of riparian and aquatic habitats and the species which inhabit them.



Fig 1.1 Carlingford area looking east from the Newry River estuary towards the mouth of Carlingford Lough





Fig 1.11 Carlingford area looking west from the Newry River estuary towards Newry




Fig 1.12 Foyle and Carlingford catchments illustrating some of the main tributaries.




Fig 1.13 Carlingford area and tributaries.

## 1.2 Atlantic Salmon and Sea Trout

Salmon and Sea Trout are referred to as being anadromous meaning that they migrate between the freshwater and marine environments returning to freshwater to reproduce. This complex life history exposes them to varied environmental pressures and recreational and commercial fisheries. Adult Atlantic salmon return to their natal rivers where spawning takes place. Sea trout also demonstrate an ability to return to their natal river but their homing instinct may not be as strong as those of the Atlantic salmon. After the eggs hatch the juveniles (initially referred to as fry and then parr) remain in freshwater for up to three years.

Smoltification is the physiological adaptation, which occurs when the juvenile salmon change from the parr stage (freshwater phase) to the smolt stage (marine phase). Smoltification at the latitudes of the Foyle and Carlingford areas tends to occur after one, two or three years. Most salmon from the north of Ireland (referred to as post smolts) will remain after smoltification in the North Atlantic for one year and are referred to on their return to the coast and rivers as grilse. Salmon that stay at sea for longer than one year are referred to as multi sea winter (MSW) salmon.

## 1.3 Non Salmonid Fish Species

As highlighted earlier populations of other non-salmonid fish species occur within the Carlingford catchments. At present monitoring is targeted at salmonid and to a lesser degree coarse species however with obligations under the Water Framework Directive other non salmonid fish species are being monitored more closely. Fish species presence, abundance and age structure can act as a good environmental/ecological indicator demonstrating the ability of the aquatic habitat to support a diverse array of native species. Populations of the European Eel, Bass, Grey Mullet, River/Brook and Sea Lamprey form an important part of the native fisheries biodiversity of the Carlingford catchments. Maintaining high standards of water quality and appropriate habitat for these species is essential for the overall health of the aquatic ecosystem. In the Carlingford area significant non-native fish species have colonised heavily modified and artificial water bodies such as Newry canal and Camlough. The Loughs Agency recognises the importance of the coarse fish populations in terms of a recreational resource for both local residents and tourists and views the improvement and development of the infrastructure to sustainably exploit this resource as a core responsibility.



Fig 1.3 Bream specimens recovered during a coarse fish survey of Newry Canal in 2002

## 2.0 ATLANTIC SALMON STOCKS

In order to describe the status of salmon stocks each of the following points need to be considered:

- Redd Counts
- Juvenile abundance
- Marine survival
- Adult abundance
- Exploitation

### 2.1 Redd Counts

Redds are spawning nests created by salmon or trout. Differentiation between salmon and trout redds can be made as salmon redds tend to be larger in size and trout tend to spawn earlier than salmon. Research within the Foyle system using extensive annual redd count data has highlighted a good relationship between the number of redds and the total annual catch of salmon. Water flow is of great significance when monitoring redds as in high water conditions the ability to see and count redds in rivers is impaired. The Loughs Agency will continue to expand redd counting in the 2008/2009 spawning season on the Whitewater River and sections of the Clanrye River.

### 2.2 Juvenile Abundance

Within the Loughs Agency jurisdiction trends in abundance of juvenile salmonids are monitored by annual semi-quantitative electrofishing surveys. The numbers, age and species of fish captured during five minute timed electrofishing surveys are compared with previous years data allowing for change to be monitored, facilitating suitable fishery management practices to be implemented. In 2008 a total of 57 sites were semi-quantitatively electrofished within the Carlingford system. The results for each site for salmon and trout are classified as excellent (>25 fish), good (15-24 fish), fair (5-15 fish), poor (1-4 fish) and absent (0 fish), Table 2.2. Figures 2.22-2.25 outline the salmon 0+ electrofishing results and site classifications for the Carlingford catchments in 2008. Please consult previous status reports for site classifications in other years.

Semi-quantitative electrofishing was developed to monitor 0+ salmonids (fry/young of the year). In order to quantify the abundance of 1+ salmonids (parr and older) fully quantitative electrofishing surveys are required which can be used to calculate fish densities within a defined area. Rivers and tributaries with good environmental quality are more likely to support good populations of each year class.

Fish populations can vary considerably over time and location, it is therefore necessary to monitor the populations over a period of years to highlight meaningful



trends before considering remedial activities such as habitat improvement works. These trends are being continually monitored by the Loughs Agency and the most appropriate management options considered.

There are a variety of reasons why electrofishing sites may be perceived to be under producing, these can include, lack of suitable juvenile habitat, the presence of impassable obstacles to migratory fish species on lower sections of a tributary, pollution, inconsiderate channel maintenance, tunnelling by bank side vegetation, stream gradient and poor forestry practices etc. The critical point is to recognise the major factors at play and to investigate all possible reasons for underproduction accepting that there may be inherent reasons as to why production may not be improved upon in certain areas. When the same areas are surveyed for other non salmonid species it may be discovered that they provide habitat more suited to these species. Habitat improvement works and the rationale behind them are discussed in greater detail later. Obligations under the Water Framework Directive are driving quantitative surveys of both salmonid and non salmonid species under proposed Surveillance, Operational, Investigative and Protected Area monitoring programmes.






Symbol	Grade	Number of 0+ Salmonids
	Excellent	>25
	Good	15-24
	Fair	5-14
	Poor	1-4
	Absent	0

Table 2.2 Loughs Agency semi-quantitative electrofishing classification system for 0+ salmon and trout



Fig 2.21 Electrofishing on the Clanrye River and salmon parr




Fig 2.22 Carlingford Area fry index 2000-2008






Fig 2.23 White Water catchment fry index 2000-2008.




Fig 2.24 Clanrye River catchment fry index 2000-2008




Fig 2.25 Salmon 0+ electrofishing site classification 2008

### 3.0 TROUT STOCKS

Annual trends in the populations of juvenile trout are also monitored within the Loughs Agency jurisdiction using the same methodology and classification system as those employed for salmon. The semi quantitative electrofishing results for trout fry in the Carlingford catchments and site classifications are displayed in Figures; 3.1 to 3.13.



Fig 3 Electrofishing survey and trout parr




Fig 3.1 Carlingford Area fry index 2000-2008.






Fig 3.11 Whitewater trout fry index 2000-2008.




Fig 3.12 Clanrye trout fry index 2000-2008.




Fig 3.13 Trout 0+ electrofishing site classification 2007

## 4.0 MARINE SURVIVAL

The numbers of salmon that survive to return to the freshwater environment are greatly influenced by conditions in the marine environment. Climate change leading to changes in sea surface temperatures, prey abundance, high seas fishing, marine pollution, sub lethal levels of pollution and predation all have an effect on the Atlantic salmon and indeed other migratory fish species chances of survival.

Marine survival trends are monitored on a number of index rivers in the North East Atlantic where total trapping facilities are available for both migrating juvenile and adult populations. Total trapping allows for an accurate count of all migrant smolts (total freshwater production) and returning adults to be made and therefore an accurate estimate of marine survival. These projects are facilitated by the use of Coded Wire Tags (CWT). Coded wire tags are small (2-3mm long) micro tags that are injected automatically by a CWT device into the snout cartilage of anaesthetised fish remaining there for the duration of the life of the fish. CWT fish also have their adipose fin (small fin between the dorsal fin and caudal fin (tail fin)) removed so that they can be identified in the various fisheries that may intercept them. In Ireland a comprehensive screening programme is conducted at all major landing ports and markets. This programme is important in monitoring the effect of the remaining salmon fisheries on salmon stocks from rivers both within and outside of the island of Ireland.

Trends in marine survival for the River Bush (nearest index river to the Foyle system) confirm patterns observed elsewhere on the southern stocks of North Eastern Atlantic salmon, which indicate that marine survival can be variable between stocks and years. In the River Bush marine survival has decreased considerably over recent years as outlined in Table 4.

<b>Year of Smolt Cohort</b>	<b>Year of Returning 1SW Grilse</b>	<b>Marine Survival %</b>
<b>Pre 1996</b>	Pre 1998	Circa 30%
<b>2002</b>	2003	5.9
<b>2003</b>	2004	4.3
<b>2004</b>	2005	4.6
<b>2005</b>	2006	4.2
<b>2006</b>	2007	13.0
<b>2007</b>	2008	7.5

Table 4 Marine survival rates for the River Bush of 1SW grilse (after exploitation at sea) pre 1996 and 2002-2007 smolt cohort. Data supplied by Agri Food and Bioscience Institute, River Bush Salmon Research Station



The figures outlined in table 5 are mirrored by those for other index rivers monitoring the southern stocks of North Eastern Atlantic salmon populations. These figures suggest that salmon are facing increased pressure for survival at sea. A major new international research project called SALSEA - Merge has been developed by scientists from the North Atlantic Salmon Conservation Organisation (NASCO) parties and its research wing the International Atlantic Salmon Research Board (IASRB). There are twenty consortium members in total including the Loughs Agency. SALSEA aims to monitor how Atlantic salmon use the ocean; where they go; how they use ocean currents, and the ocean's food resources, and what factors influence migration and distribution at sea. Research cruises commenced in 2008 and will continue in 2009 to collect the necessary data to answer the questions listed above. Over 426 post smolts were caught by the two Irish cruises and 363 post smolts caught by the Faroese in the areas highlighted below. Further information and project details can be found at: <http://www.nasco.int/sas/salsea.htm>




Fig 4 Proposed marine survey areas for salmon in 2008



Fig 4a RV Celtic Explorer SALSEA research cruise 2008



Figure 4b Picture from the Irish Research Vessel Celtic Explorer taken during the second SALSEA research cruise 16-24<sup>th</sup> May 2008



Since 2003 partial smolt trapping including CWT tagging has been conducted in the Faughan catchment using a rotary screw trap, Fig 4c.



Figure 4c Rotary screw trap in position on the River Faughan directly below the fish pass at Campsie barrage.

Smolt trapping can have a number of objectives including the monitoring of both salmonid and non salmonid species. Sampling of the age composition, obtaining information on run timing and recording length/weight data is conducted in tandem with the tagging programme. As mentioned above total counts of migrating smolts can be made on rivers. Where this is unfeasible due to the absence of total trapping facilities, total smolt migration can be estimated by means of a mark-recapture experiment.

In 2004 an estimate of total smolt production for the Faughan catchment was made by a mark-recapture study resulting in a minimum run size estimate of 33,854 migrating salmon smolts. The estimate was a minimum due to a number of high water events that prevented the smolt trap from fishing for a period of time during the peak smolt migration period. Tables 4.1 and 4.12 outline numbers of salmon smolts tagged from 2003-2008 and recapture data for 2004 and 2007.

Year	No of Salmon Smolts Tagged	Average Length (mm)	Average Weight (g)
2003	2113	149	33.45
2004	2500	134	24.6
2005	2210	133	23.6
2006	1025	133	25.36
2007	2062	135	27.1
2008	1865	130	22.1

Table 4.1 Numbers and average weight and length of salmon smolts tagged on the River Faughan 2003-2008. Coded Wire Tagging equipment was purchased by the Loughs Agency in 2005 with funding secured from the European Regional Development Fund through the INTERREG IIIA Programme, administered by the Environment and Heritage Service, on behalf of the Department of Environment.

Year Tagged	Year Recaptured	Numbers Recaptured	Recapture Location
2003	2004	12	Greencastle, Burtonport, Malin Head, Belmullet and Torr Head
2004	2005	16	Greencastle, Malin Head, Donegal and Galway Bay
2005	2006	3	Greencastle
2006	2007	2	Greencastle and Ballycastle

Table 4.12 Recapture data from River Faughan CWT programme. Data for fish tagged in 2007 and recovered in 2008 will not be available until 2009.




Figure 4d Salmon smolt run timing and abundance from rotary screw trap sub sample, River Faughan 2004-2008. Breaks in data are due to closure of trap during high water conditions

In 2004 a detailed examination was carried out on the age class of migrating salmon smolts in the Faughan catchment, Table 4.13.

Age at Smolting	%
1	13
2	83
3	4

Table 4.13 Age class of salmon smolts migrating from the Faughan catchment in 2004



Fig 4.14. From top to bottom, Atlantic salmon smolts from the Faughan Catchment, brook lamprey, river lamprey and sea lamprey also caught in the River Faughan smolt trap

## 5.0 ADULT ABUNDANCE

Adult Atlantic salmon abundance is assessed in three ways: directly by using commercial netting/recreational rod catches and fish counters and indirectly by reference to conservation limits/spawning targets.

Using catch data as a measure of population status is a well established and extensively used technique. In the Foyle system annual commercial and recreational catch data has been recorded since the establishment of the Foyle Fisheries Commission in 1952, with some data available before this period. Within the Carlingford area catch data is available from 2001. No commercial salmonid fisheries are pursued in the Carlingford system. The relationship between catch and stock is complex and care should be applied in interpretation. A more precise measure of catch incorporates fishing effort (number of licences issued or the amount of time fished) and is referred to as catch per unit effort (CPUE).

### 5.1 Recreational Fisheries

One problem encountered when analysing catch data is unreported catch. All recreational fishers are required by law to make catch returns. This information facilitates management decision making and therefore it is vitally important that all catch returns are accurate and made promptly at the seasons end.

Year	Declared Rod Catch Salmon	Declared Rod Catch Sea Trout	Returns as a % of Licences Issued
1999	1022	679	3.74
2000	723	417	2.55
2001	3188	450	17.68
2002	5117	1010	27.93
2003	1844	361	15.5
2004	2285	75	13.99
2005	4084	413	25.77
2006	3476	469	37
2007	4929	379	22.11
2008	4060	815	54.94

Table 5 Declared rod catch returns for salmon and trout in the Foyle and Carlingford areas. Note figures include the Clanrye and Whitewater in the Carlingford area from 2001 onwards. Carcass tagging was introduced in 2001.



Year	Declared Catch Carlingford System (Salmon)	Declared Catch Carlingford System (Trout)
2003	0	0
2004	17	3
2005	0	33
2006	3	8
2007	44	46
2008	62	45

Table 5.1 Declared catch from the Carlingford system for salmon and trout 2001-2008



Fig 5.11 Angler on the upper reaches of the Faughan River

## 5.2 Commercial Fisheries

Commercial fisheries have traditionally operated within the Foyle sea area, Lough Foyle and tidal River Foyle. As mentioned above no commercial fisheries for salmon are pursued within the Carlingford system. Within the Foyle area the drift net and draft net fisheries as well as the rod fisheries have been closely regulated with a real time management regime in place to monitor the numbers of fish migrating up key rivers. If predetermined numbers of fish have not been counted by the strategically placed electronic fish counters at Sion Mills weir (River Mourne), Campsie Barrage (River Faughan) and the Plumb Hole (River Roe) then specified closures of the commercial and/or recreational fisheries are enforced.

In 2007 new regulations were introduced to reduce the number of commercial nets operating within the Foyle area and all mixed stock interceptory drift nets seaward of Lough Foyle were curtailed. This decision was made to comply with the EU Habitats Directive, similar curtailment of mixed stock fisheries were introduced in the Republic of Ireland. Within the Foyle area this was achieved through a voluntary hardship scheme. 18 out of 112 drift nets remain in Lough Foyle, those remaining have been reduced in size from 900m to 500m and 10 out of 50 draft nets remain. This represents a significant reduction of netting effort. Regulations were also introduced to limit the numbers of fish which could be retained by the recreational rod fishery throughout the Foyle and Carlingford areas.

Year	Drift Catch	Draft Catch	Total Drift and Draft
1998	31296	11141	42437
1999	15397	7893	23290
2000	22333	10339	32672
2001	13500	9476	22976
2002	28851	11917	40768
2003	15741	16991	32732
2004	12800	9490	22290
2005	13391	12143	25534
2006	6160	6031	12191
*2007	2598	2774	5372
2008	1248	2924	4172

Table 5.2 Declared catch from the commercial salmon fisheries 1998-2008. Note 100% rate of catch returns. \* Reduced numbers of commercial nets operating in the Foyle area from 2007



Fig 5.21 Commercial Fishing. Draft netting on the tidal River Foyle and drift netting in Lough Foyle

### 5.3 Counters

Within the Foyle system a number of river catchments have electronic fish counting facilities that provide estimates on the run timing and abundance of fish >45cm. A Logie resistivity fish counter has been installed on the Newry River within the Carlingford system (figure 5.3), this facility has been operational since September 2007. The new counting facilities will provide valuable information on the run timing and abundance of fish in the Newry/Clanrye River and will facilitate future fisheries management decision making. Counts for the Newry River catchment within the Carlingford area are outlined in table 5.3 and figure 5.31.



Fig 5.3 Fish Pass and counter construction on the Newry River.

Year	Number of fish >45cm
*2007	*32
2008	268

Table 5.3 Newry/Clanrye River fish counter figures. \*Note the Newry/Clanrye fish counter was installed in September 2007. The figures reported are indicative and form the initial output from the commissioning phase of the counter installation.




Fig 5.31 Monthly fish counts on the Newry River Sep 2007-2008

## 5.2 Conservation Limits/Spawning targets

Another way to assess adult salmon stock status is to monitor run sizes on rivers and to compare them with predefined reference points called conservation limits. Conservation limits define a level of spawning that optimises the sustainable catch by commercial and recreational fisheries. If exploitation rates increase above the sustainable catch levels the catch may increase in the short-term but the stock will eventually reduce. Conservation limits demarcate the undesirable spawning stock level at which recruitment would begin to decline significantly (NASCO). The real time management regime incorporating the setting of management targets and spawning targets implemented in the Foyle aims to manage the fisheries and spawning populations in a sustainable manner. The management and spawning targets are set for the various river catchments based on the amount and quality of nursery habitat present. River habitat surveys are carried out along each stretch of river and graded according to the type and quality. Egg deposition levels are set according to the quality grading of each section of nursery habitat.

There are four grades of nursery habitat, however for the purpose of setting egg deposition levels only grades 1-3 are utilised. Grade 1 denotes the best quality habitat. The egg deposition rate/carrying capacity is set as follows. Grade 1 = 10 eggs per m<sup>2</sup>, grade 2 = 5 eggs per m<sup>2</sup> and grade 3 2.5 eggs per m<sup>2</sup>. The total number of eggs is calculated by multiplying the area of each grade of nursery habitat by the appropriate density of eggs per m<sup>2</sup>. 25% is deducted from the management target

allowing for loss of salmon by angling (15%) and poaching and predation (10%). The remaining figure is referred to as the conservation limit/spawning target.

Once the number of eggs required for each river has been established this can be converted to a total number of fish required to achieve the management targets and conservation limit/spawning targets. The average fecundity (number of eggs produced per female) of Foyle salmon has been estimated at 2500 and the ratio of female to male salmon estimated at 60:40. When combined with the amount of nursery habitat of the various grades this equates to the conservation limit/spawning target.

In the Carlingford area targets can be set and monitored once adequate information has been collated from both fish counting facilities and recreational catch returns from the salmon fisheries. Where necessary catchments which do not meet their targets may have fishery conservation measures imposed and or fish stock rebuilding programmes instigated.

## 6.0 COARSE FISH STOCKS

Coarse fish species differ significantly from most salmonid species in that their lifecycle is completed solely in freshwater. Coarse species also utilise differing habitat types than salmonid species preferring slower moving deeper water. Newry canal and Camlough are good examples of coarse fish habitat within the Carlingford area. The coarse fish species present within the Carlingford area are not native to the island of Ireland but have been introduced over the last few hundred years as a source of food, for sport and by escaping from fish farms. Irelands natural water courses are defined primarily as salmonid waters however artificial water bodies such as canals and reservoirs provide ideal habitat for a variety of coarse fish species. These artificial or heavily modified water bodies facilitated the colonisation by coarse fish species providing suitable habitat for spawning, nursery areas for juveniles and appropriate conditions and feeding opportunities for adult populations.

The Loughs Agency aims to provide sustainable social, economic and environmental benefits through the conservation, protection and development of the coarse fisheries of the Foyle and Carlingford areas by promoting the sustainable exploitation of the resource to achieve maximum benefit to local communities.

In 2008 the Loughs Agency in collaboration with the Agri Food and Bioscience Institute (AFBI) conducted a survey of Derryleckagh Lough which discharges into the Clanrye River. The aims of the survey were to chart the bathymetry/depth of the lough and to identify the fish species present.



Derryleckagh Lough is a 30 hectare eutrophic lake situated in County Down, Northern Ireland. It has a maximum depth of 9.0m and an average depth of 2.0m, with much of the southern end of the lake being <2m deep. The lake contains pike and eels, and one small perch specimen was captured in this survey.

At the time of the survey, water temperatures did not differ greatly among depths with a difference of <2°C between the surface and 7m. All depth strata within the lake had sufficient dissolved oxygen to support fish, ranging from 97.9% at the surface to 73.9% at 7m (Figure DL 1). Total Phosphorous recorded at the time of the survey was 59µg l<sup>-1</sup>, classifying the lake as eutrophic.




Figure DL 1. Temperature/DO profile in Derryleckagh Lough

This brief report provides a summary of the fish survey on Derryleckagh Lough, undertaken from 28-29<sup>th</sup> August 2008, including species, number, length frequency, age and locations of fish captured.

**Lake:** Derryleckagh Lough

**Irish Grid Reference:** J128 256

**Survey date:** 28-29.08.08

**Weather:** Cloudy, dry, light SSE wind

**Air temperature:** 18°C

**Surface water temperature:** 17.1°C

**Bank type:** Stones, reeds, grass, overhanging trees

**Net types (number):**

Lundgrens Norden, 30m, 12 panel multimesh monofilament gill nets (8)

Lundgrens, 30m, 75mm fixed mesh monofilament gill net (1)

Lundgrens, 30m, 60mm fixed mesh monofilament gill net (1)

50cm x 10m fyke nets, chains of 3 (3)

## Results and Discussion

A total of only 12 fish were captured during this survey; 8 pike, 3 eels and 1 perch. Net locations and the number of fish caught in each net are summarised in Figure 2 and Table DL 1 below. Although few fish were captured overall, it might have been expected to capture more pike in the shallower nets than the deeper set nets. This proved not to be the case however, with most of the pike being captured in nets >3m deep. Pike ranged in size from 16 to 42cm (Figure 3) and age from 0+ to 2+ years. One eel was captured in each of the three fyke nets, ranging in size from 41 to 55cm (Figure 4). Only one small perch was captured, 5.7cm in length. With small perch being a shoaling fish, this is very surprising. The possibility exists that the fish was captured in a previously surveyed lake and was already present in the gillnet before it was set, however this is unlikely as it would be easily seen whilst setting. It could be, however, without the presence of perch as a prey species, that the lake contains a population of mainly cannibalistic pike with a small number of large individuals preying on, and to an extent 'controlling', the number of smaller pike in the population. Such large pike would not likely be captured using the monofilament gillnets adopted for this survey.

A concurrent hydro-acoustic survey conducted at night-time showed very few fish tracks in general, and none in the open water pelagic zone. This would support the idea of a lake containing only pike (and eels), as shoals of perch generally disperse at night-time and are easily detected using horizontal hydro-acoustics if present.

Further surveys would be required to ascertain the presence/absence or size of any perch population within the lake; however evidence from this survey certainly suggests that if a population does exist it is likely to be very small.




Figure DL 2. Location of benthic gillnets (red circles), floating gillnet (white circle) and fyke nets (red squares) in Derryleckagh Lough

Net (type)	Mesh size (mm)	Depth	Perch	Eels	Pike	Total
G1 (Norden 12 panel gill)	5 - 55	0-3m	1			1
G2 (Norden 12 panel gill)	5 - 55	0-3m			1	1
G3 (75mm fixed mesh gill)	75	0-3m				
G4 (60mm fixed mesh gill)	60	3-6m				
G5 (Norden 12 panel gill)	5 - 55	6-12m			2	2
G6 (Norden 12 panel gill)	5 - 55	6-12m				
G7 (Norden 12 panel gill)	5 - 55	3-6m			4	4
G8 (Norden 12 panel gill)	5 - 55	3-6m				
G9 (Norden 12 panel gill)	5 - 55	3-6m			1	1
G10 (Norden 12 panel gill)	5 - 55	Floating				
F1 (fyke)		0-3m		1		1
F2 (fyke)		0-3m		1		1
F3 (fyke)		0-3m		1		1
			1	3	8	12

Table DL 1. Number of fish caught in each net in Derryleckagh Lough




Figure DL 3. Length frequency of pike in Derryleckagh Lough




Figure DL 4. Length frequency of eels in Derryleckagh Lough





Figure DL 5. Derryleckagh Lough survey 2008

In 2007 a resurvey of the coarse fish populations of the Newry Canal was conducted by the Loughs Agency in collaboration with the Agri Food and Biosciences Institute and the Central Fisheries Board. The 2007 survey consisted of both netting and horizontal hydro-acoustic techniques during day time and night time hours. The hydro-acoustic method results in fewer gill nets being set and records data passively. The full report on the Newry Canal Fish Stock Assessment can be downloaded from the Loughs Agency website [www.loughs-agency.org](http://www.loughs-agency.org). The canal was surveyed from the Albert Basin to Victoria Lock. Table 6 outlines summary results from the 2007 survey.

Species	Number Caught	Length Range (cm)
Roach	144	8-26
Bream	1	21
Roach x Bream	16	8-30
Pike	5	44-75
Perch	36	9.7-27
Tench	4	26

Table 6 Fish survey results from Newry canal 2007. Eel and Flounder were also recorded.



Fig 6 Newry Canal fish stock assessment 2007

In 2001 and 2002 the Loughs Agency in collaboration with the Central Fisheries Board conducted baseline surveys of the fish species present within Newry canal between the town of Newry and Victoria Lock. This was designed to assess the population status of the fish stocks. Gill and fyke nets were used to capture fish with a proportion of all fish being measured, weighed and scaled for subsequent age analysis. Tables 6.1 and 6.2

Species	Number Caught	Length Range (cm)	Weight Range (kg)
Pike	118	13-96	0.03-7.5
Roach	207	12.5-25	0.1-0.3
Bream	26	15-38	0.05-0.9
Roach x Bream	9	26-29	Average 0.95
Tench	2	34 & 37.5	0.9 & 0.95
Eels	>250	N/A	>0.5-1.4
Brown Trout	1	23.5	0.2

Table 6.1 Fish survey results from Newry canal 2001

Species	Number Caught	Length Range (cm)	Weight Range (kg)
Roach	437	5-27	Up to 0.43
Bream	36	20-46	Up to 1.8
Roach x Bream	58	26-35	0.35-0.85
Pike	40	35-73	Up to 3.45

Table 6.2 Fish survey results from Newry canal 2002. Perch, Brown trout and Eels were also caught in 2002







Fig 6.1 Left to right Perch, Pike and Bream

## 7.0 HABITAT MONITORING

The Loughs Agency has carried out extensive habitat surveys on all the major rivers and tributaries within the Foyle and Carlingford catchments. Habitat surveys are carried out on foot. Although time consuming this is at present the best method for classifying the various grades of habitat. Habitat is classified into one of three life cycle units Fig 6, the presence and order of which is essential to the productive capacity of a salmonid river. Other non salmonid species also benefit from diverse in-channel habitat. The life cycle unit categories include spawning, nursery and holding habitat. Each category is then graded on a scale of 1-4, 1 representing the best quality attainable and 4 the worst. Other data collected during these surveys include channel width and impassable barriers to migratory fish species.




Fig 7 Life cycle unit depicting the type of habitat found in spawning, nursery and holding zones



Fig 7.1 Examples of spawning, nursery and holding habitat



## 8.0 LAND USE

Land use classification is an important tool when assessing the potential impacts within a particular river catchment or indeed when looking at specific land use and land management practices. Land use impacts could have either a positive or negative impact on rivers and tributaries. A good understanding of the land use within a catchment is therefore imperative in managing at a catchment scale.

Land use in Northern Ireland has been captured using satellite imaging technology and classified to type. The following figures outline the broad land use classification within the Carlingford Area.




Fig 8 Carlingford Area land use classification




Fig 8.1 Carlingford Area land use classification map



## 9.0 WATER QUALITY

Routine water quality monitoring within the Foyle and Carlingford areas is conducted by the Northern Ireland Environment Agency (NIEA) of the Department of the Environment for Northern Ireland and the County Councils in the Republic of Ireland (Donegal County Council and Louth County Council). Routine sampling is conducted regularly for both chemical and biological General Quality Assessments (GQA).

In addition to the routine river monitoring carried out by the NIEA and the County Councils the Loughs Agency conducts proactive and reactive pollution investigations to investigate or highlight problems or potential problems which may have an effect on the aquatic environment and ultimately on the fish species and aquatic habitats.

In 2008 the Loughs Agency continued a programme of monitoring at the tributary level for assessments of chemical and biological water quality which was instigated in 2007. Tributaries within the Carlingford area are monitored for chemical water quality parameters including Biological Oxygen Demand (BOD), Suspended Solids, Ammonia and Phosphorous. Biological water quality was assessed using the Biological Monitoring Working Party (BMWP) a biotic scoring index.



Fig 9.0 Loughs Agency chemical water quality testing in the laboratory

The Loughs Agency also maintains a mobile pollution response unit containing aerating equipment and absorbent and non absorbent booms for oil and chemical spills. The unit can be rapidly deployed to the site of a pollution incident.

### **Water Quality Parameters**

The following water quality parameters are monitored through the Loughs Agency monitoring programme and determined from water samples in the laboratory:

- Biochemical Oxygen Demand (BOD)
- Ammonia
- Phosphorus
- Suspended Solids

### **BOD**

Any organic matter discharged into a river provides an immediate source of food for bacteria. These bacteria will break down the organic matter eventually into simple compounds such as carbon dioxide and water. Biochemical Oxygen Demand or Biological Oxygen Demand (BOD) is a chemical procedure for determining how fast biological organisms use up oxygen in a body of water. It is considered as an indication of the quality of a watercourse

### **Ammonia (NH<sub>3</sub>)**

Ammonia is generally found in small amounts in rivers and streams. This is due to microbiological activity and the resultant reduction of compounds containing nitrogen. High levels of ammonia can occur as a result from sewage pollution and have detrimental impacts on fish species.

### **Phosphorus (PO<sub>4</sub>)**

The over-loading of nutrients such as phosphorus in watercourses often leads to a process known as eutrophication. Eutrophication is a major environmental issue in Irish rivers and lakes. Sources of phosphorus include agricultural fertilizers and household detergents.

### **Suspended Solids**



Particulate matter may be organic or inorganic in nature. Organic solids may consist of algal growths, indicative of eutrophic conditions. Inorganic solids generally are the result of discharge washings from sand and gravel extraction activities or quarries. Suspended solids can affect plant growth and fish habitats.

The following parameters are also recorded at each sample station by means of an electronic measuring probe:

- pH
- Temperature
- Dissolved Oxygen
- Conductivity

## **pH**

This is a measure of the hydrogen ion concentration of a solution and therefore an indication of whether a liquid is acid or alkaline. The pH scale ranges from 0 (very acid) to 14 (very alkaline), with results generally influenced by geological conditions. Fish can be susceptible to changes in pH. Low pH levels are generally found in catchments with high forestry operation impacts.

## **Temperature**

The effect of changes in temperature on living organisms, such as fish, can be critical. Thermal discharges from urban and industrial sources can lead to temperature increases in watercourses and increased stress on aquatic habitats and associated species.

## **Dissolved Oxygen**

Sufficient levels of oxygen saturation in fresh waters are generally an indication of good ecological status and ideal for fish life. The main point to remember about oxygen solubility is that it has an inverse relationship with temperature. This helps explain why DO levels are generally lowest during summer low flow conditions, increasing the risk of pollution from discharges at this time.

## **Conductivity**

The conductivity or electrical conductivity of a watercourse is a measure of its ability to conduct an electric current. Electrical conductivity estimates the amount of total dissolved salts, or the total amount of dissolved ions in the water. Electrical Conductivity is controlled by geology and any variations may be sourced to increased ions from wastewater from sewage treatment plants or urban run-off from roads.




Fig 9.1 Carlingford area average suspended solids results 2008. Values are in mg/l






Fig 9.2 Carlingford area Ammonia results 2008. Values are in mg/l






Fig 9.3 Carlingford area phosphorous results 2008. Values are in mg/l






Fig 9.4 Carlingford area Biological Oxygen Demand (BOD) results 2008. Values are in mg/l






Fig 9.5 Carlingford area Biological Monitoring Working Party results 2008

NIEA routinely monitor both the chemical and biological water quality within the rivers of Northern Ireland. In relation to chemical monitoring an extensive network of sampling stations are monitored for a variety of chemicals. The General Quality Assessment (GQA) is defined by limits for the concentrations of Biological Oxygen Demand (BOD), ammonia and dissolved oxygen (DO). The measures listed are indicators of the affect on water quality by waste water discharges and agricultural run-off containing organic material. Water quality can be affected by a variety of sources and the GQA determinands provide a recognised assessment of water quality.

The overall GQA class assigned to a section of river is based on the worst performing of the three measures (BOD, ammonia and DO). Table 9.1 outlines the standards for the chemical GQA.

<b>GQA Class</b>	<b>Dissolved Oxygen (% Sat) 10- percentile</b>	<b>BOD (mg/l) 90-percentile</b>	<b>Ammonia (mg/l) 90-percentile</b>
<b>A (Very Good)</b>	80	2.5	0.25
<b>B (Good)</b>	70	4	0.6
<b>C (Fairly Good)</b>	60	6	1.3
<b>D (Fair)</b>	50	8	2.5
<b>E (Poor)</b>	20	15	9.0
<b>F (Bad)</b>	<20	-	-

Table 9.1 chemical GQA class limiting criteria

The above table can be summarised as follows: for BOD and ammonia the section of river should contain less than the stated levels for at least 90% of the time. DO levels must not fall below the stated levels for more than 10 percent of the time.

In relation to biological monitoring an extensive network of sampling stations is also routinely monitored. The biological GQA is defined by observed measures of the abundance and diversity of macro invertebrates (for example freshwater shrimps, insect larvae and molluscs) compared to expected values as derived from a UK computer model adapted for Northern Ireland called River Invertebrate Prediction and Classification System (RIVPACS)

Different species of macro invertebrates are more sensitive to specific forms of pollution and therefore environmental quality indices (EQIs) based on biological results may be used to assess water quality. Macro invertebrates are also the dominant prey of both salmonid and some non salmonid fish species. The measure of diversity of a macro invertebrate community can be a more reliable indicator of the pollution pressures within a catchment than relying solely on an assessment of chemical water quality. The impacts of pollution on a macro invertebrate community are longer lasting and can highlight intermittent pollution impacts that may be missed through chemical water quality monitoring.



Biotic scoring systems have been developed to assign a score based on a standardised system to each sample site. One such system is the Biological Monitoring Working Party (BMWP). Generally the higher the BMWP score the better quality of the macro invertebrate community which reflects better water quality.

Based on a combination of biotic scoring systems biological GQA classes are assigned to sections of river. The two EQIs used are as follows:

$$\text{EQI}_{\text{taxa}} = \frac{\text{BMWP Observed Number of Taxa}}{\text{BMWP Predicted Number of Taxa (as derived from UK model)}}$$

BMWP Predicted Number of Taxa (as derived from UK model)

$$\text{EQI}_{\text{ASPT}} = \frac{\text{BMWP Observed ASPT (Average Score Per Taxon)}}{\text{BMWP Predicted ASPT (as derived from UK model)}}$$

BMWP Predicted ASPT (as derived from UK model)

Biological Class	EQI for ASPT	EQI for Taxon
<b>A (Very Good)</b>	1.00 or above	0.85 or above
<b>B (Good)</b>	0.90-0.99	0.70-0.84
<b>C (Fairly Good)</b>	0.77-0.89	0.55-0.69
<b>DC (Fair)</b>	0.65-0.76	0.45-0.54
<b>E (Poor)</b>	0.50-0.64	0.30-0.44
<b>F (Bad)</b>	<0.50	<0.30

Table 9.2 Biological GQA class limiting criteria



Figure 9.6 Flattened mayfly nymph from the order *ephemeroptera* high scoring macro invertebrate indicative of good water quality

European Council Directive 92/43/EEC of the 21st of May 1992 on the Conservation of Natural Habitats and on Wild Flora and Fauna (Also known as the Habitats Directive) was enacted in Northern Ireland under the European Communities Nature Conservation (Natural Habitats etc.) Regulations (Northern Ireland) 1995.

This indicates that those areas designated as areas of nature conservation designated for salmon should strive to achieve the water quality targets that are necessary for the designated species, which has additions to the GQA standards.

While it is current government policy for all rivers to meet the General Quality Assessment Standards, the Agency feels that favourable conditions standards as detailed below should be the water quality targets for all salmonid rivers within its jurisdiction.

### 9.1 Favourable condition tables, target levels

Natural Heritage of Northern Ireland Environment Agency have suggested guidelines for the determination of water quality, the first being the proposed UK Guidance on Conservation Objectives from monitoring designated sites and includes the following, which are considered as the favourable conditions tables.

They recommend Biological GQA Class A or B with no drop in class from the existing station, and Chemical GQA Class A or B depending on which type. This is in addition to no drop in class from the existing station. In addition to these favourable conditions tables, based on publications from Conserving Natura 2000 Rivers, the European Life Series, Ecology Series; No 7 Ecology of the Atlantic Salmon, *Salmo Salar* L. these publications have indicated that there are specific favourable conditions for this species.

An annual mean of less than 10 milligrams per litre suspended solids for nursery grounds, and annual mean of less than 25 milligrams per litre for migratory passage and the setting of soluble reactive phosphorous targets in relation to river reach types which should be as near background levels.

Parameter	Level	Percentile	Reason
<b>BOD mg/l</b>	2.5	90	GQA class A
<b>Ammonia mg/l</b>	0.25	90	GQA class A
<b>Dissolved Oxygen % Saturation</b>	80	10	GQA class A
<b>Unionised Ammonia mg/l</b>	0.025	95	Favourable Conditions Habitat Forming
<b>Suspended solids mg/l</b>			Specific for Atlantic Salmon
Nursery grounds	10		
Migratory passage	25		
<b>Soluble Reactive Phosphorous mg/l</b>	Background	-	Specific for Atlantic Salmon

Table 9.1 Favourable condition targets for Atlantic salmon

The Water Quality data in the reports has come from the Northern Ireland Environment Agency, Water Management Units Water Quality Archive. It is accepted by the Agency that monitoring is designed to ensure that water quality is monitored to ensure compliance with European Union directives. The monitoring however does not tie in well with the habitat and electrofishing survey monitoring carried out by the Loughs Agency, and as such the Loughs Agency instigated its own monitoring programmes in 2007 to link fish life, macro invertebrates and water quality into one holistic site evaluation. Additionally Northern Ireland Environment Agency, Water Management Unit data is not released in real-time and the data displayed below is for 2007, where the Loughs Agency status report is for 2008. By collecting and analysing water quality data the Loughs Agency can react to local water quality issues more effectively.

From 2008 water quality information from NIEA will be reported in a different manner. The European Union Water Framework Directive requires that numerous water quality parameters in addition to those traditionally monitored should be combined to provide an overall classification for water bodies. Additional parameters will include fish, diatoms and morphology amongst others. The Loughs Agency will monitor fish for WFD purposes and provide this information to NIEA.

The Loughs Agency will report on WFD classifications in the 2009 catchment status reports.




Fig 9.07 Chemical General Quality Assessment (GQA) Carlingford area 2007. Data supplied by NIEA






Fig 9.08 Biological General Quality Assessment (GQA) Carlingford area 2007. Data supplied by NIEA

## 10.0 CONSERVATION AND PROTECTION

The Loughs Agency continues to carry out an active fishery protection role throughout the catchments of the Carlingford area including the sea area and on all tributaries. Tables 10 and 10.1 outline the number of patrols and some duties carried out by the Loughs Agency staff in the Carlingford area and seizures for the Foyle and Carlingford areas.

A team of Fishery Officers based in Carlingford are responsible for the catchments within the Carlingford area. In addition to fishery protection duties the team is responsible for conducting sampling within Carlingford Lough.

Year	No of Licence Checks	Joint Patrols	On-site Inspections
2005	786	3	230
2006	550	4	267
2007	410	0	110
2008	447	0	37

Table 10. Breakdown of conservation and protection duties in the Carlingford area catchment 2002-2008. Note policy changed in 2008 in relation to on site inspections.

Year	2008	2007	2006	2005	2004	2003	2002
<b>Nets</b>	114	100	97	114	181	198	207
<b>Salmon</b>	92	56	91	118	130	155	94
<b>Rod &amp; Reel</b>	136	85	26	10	16	12	22
<b>Vehicles</b>	1	0	2	1	1	0	0

Table 10.1 Seized nets, salmon, rod/reels and vehicles in the Foyle and Carlingford areas 2002-2008

Year	Nets	Rod/Reel
2008	2	23
2007	2	12

Table 10.2 Seizures in the Carlingford area 2008

## 10.1 Habitat Improvement Works

In addition to the traditional protection duties carried out by the Loughs Agency staff conservation and improvement of habitat has been increasing over recent years.

Over time man has imposed significant changes on the natural courses of many rivers and flood plains. The driving forces behind these changes have included amongst others; arterial drainage schemes to provide more suitable land for agricultural purposes, urban sprawl, infrastructure expansion (roads etc.), flood defences, water abstraction and hydro power generation. All have had a significant impact on the natural meanderings and discharges of rivers and tributaries resulting in faster runoff of floodwaters ultimately leading to a change in the morphology and flow regime of rivers and resultant impacts on fisheries.

While all these processes have had some impact within the Foyle and Carlingford systems, they are still considered to be relatively natural systems with natural river structure present in the catchments headwaters. In areas that have been altered methods for reinstating lost habitat are investigated and where appropriate action taken.

In July/August 2008 the Loughs Agency, in conjunction with the Kilbroney Angling Club co-funded a project on the Kilbroney River in the Fairy Glen area. Surveying had identified that there was a lack of holding water between the mouth of the river and the top of the Glen and many of the stone weirs that were in place were in need of repair.

In-stream works were carried out to address these issues and reinstate/create pools and some spawning habitat. The works were impacted by a serious flood event which occurred in mid August but the damage was repaired by club members at their own expense.



Fig 10.4 Reinstatement of pool with some repair work to a weir on the Kilbroney





Fig 10.5 Creation of spawning habitat on the Kilbroney



Fig 10.6 Repair of weirs on the Kilbroney



Fig 10.7 Repair of damaged weir on the Kilbroney





Fig 10.8 Trimming of overhead cover and pool reinstatement on the Kilbroney



Fig 10.8 Grading of bank on the Kilbroney







Fig 10.9 Pool excavation and repair of weir on the Kilbroney



Fig 10.10 Repair of weir on the Kilbroney



Fig 10.11 Repair of weir and reinstatement of pool on the Kilbroney

## 11.0 ENVIRONMENTAL ISSUES

Some environmental issues affecting water quality have already been outlined previously. The following list presents some of the main pressures on fish populations within the Carlingford area:

- Agricultural activities – enrichment from natural and artificial fertilisers often make their way into watercourses, enhancing problems with eutrophication.
- Forestry activities – planting and felling operations can lead to increased loading of suspended solids in watercourses. Established forestry as a major upland land use has been attributed to increased acidification.
- Barriers to migration – a range of natural and anthropogenic features on rivers can lead to barriers for migrating salmonids and other fish species. These can include weirs and hydro-electric schemes.
- Gravel removal – gravel is extremely important for the creation of redds for spawning fish. Removal of gravel from the river bed in sensitive areas can destroy potential spawning and nursery habitat.
- Quarrying activities – the extraction of aggregates such as rock, sand and gravel has the potential to cause increased levels of suspended solids in nearby watercourses. Sufficient mitigation measures should be in place at such sites to trap increased sediment loads entering rivers and streams.
- Abstraction – water abstraction from watercourses for a range of uses is increasing throughout the Foyle and Carlingford catchments. Unless appropriately assessed and licenced, these activities have the potential to reduce residual flow levels and alter the ecological status of our rivers. This is even more concerning in the light of climate change.
- Peat harvesting – Peat harvesting still occurs in small upland pockets throughout the Foyle and Carlingford areas. It has the potential to increase sediment loading in receiving waters.
- Sewage treatment – sewage and waste water treatment works are under considerable pressure with the increase in urban development in our towns and villages. Several inadequate systems throughout the Foyle and Carlingford areas continue to pollute rivers.
- Hydropower – small-scale hydropower schemes are beginning to appear on rivers throughout the Foyle and Carlingford catchments. Baseline fishery data must be provided to allow for sufficient assessment of any proposed scheme, unless located above an impassable fish barrier.
- Urban development – the expansion of large-scale housing developments and the associated pressures on waste water and sewage treatment works are a potential source of water pollution in the event of overflows.
- Drainage and canalisation – these have direct impacts on the quality of available fishery habitat within the catchments. Canalisation in particular can lead to the removal of important spawning, nursery or holding areas of rivers.
- Industrial discharges – larger urban areas with industrial discharges have the potential to cause pollution through toxic discharges and can alter the temperature of the watercourse.
- Septic tanks – a proliferation of single dwellings and their septic tanks is an ongoing area of concern. Initial research from parts of the Foyle system

indicates that this is major contributor to decreased water quality and local increases in suspended solids.

## 12.0 DESIGNATED AREAS

The European Commission Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (EU Habitats Directive 92/43/EEC) requires that all member states designate Special Areas of Conservation (SACs) in order to protect threatened habitats and species. The European Commission Directive on the Conservation of Wild Birds (Birds Directive 79/409/EEC) also requires the designation of Special Protected Areas (SPA's). Together the designated SAC's and SPA's create the NATURA 2000 network of protected sites. A number of rivers have been designated as SAC's both in Northern Ireland and in the Republic of Ireland however no rivers within the Carlingford area have been designated.

NATURA 2000 sites within the Carlingford area include Derryleckagh SAC, Rostrevor Wood SAC, Slieve Gullion SAC, Carlingford Mountain SAC, Carlingford Shore SAC and Carlingford Lough SPA.

The Mourne Area of Outstanding Natural Beauty lies partially within the Carlingford area as does the Ring of Gullion AONB. The area is also covered by a variety of nationally designated sites including Areas of Special Scientific Interest (ASSI) and National Nature Reserves (NNR).

Designated sites are required to attain high environmental quality standards set at both the European and national scale. The maintenance of a network of sites represents great opportunities for co-ordinated environmental management with issues such as water quality and habitat conservation having a direct link to the quality of the fisheries resource.

## 13.0 GENETIC STUDY

A baseline genetic survey of the Atlantic salmon population was carried out in the Whitewater catchment in 2008. The results from a similar survey conducted in the Foyle area confirmed the existence of genetically distinct populations between and within the rivers and tributaries of the Foyle area. An understanding of these genetically differentiated populations is required to facilitate appropriate management of conservation measures and the commercial/recreational fisheries.

The report concluded that genetic diversity is high between and within the various salmon populations present in the Foyle system. Each population has evolved over time creating distinct populations (with some gene flow from straying fish) that are best suited to the conditions present in a particular river or tributary. The non-uniform



nature of the populations adds to the diversity of life history strategies exercised by Foyle salmon. Distinct differences such as run-timing and age at smolting can act as nature's insurance policy to any catastrophic events which would threaten a homogenous population.

The report stated that the current genetic structure and diversity of Foyle salmon is representative of what might be regarded as the native structure of wild salmon populations. The maintenance of genetic diversity is a core requirement for the long-term sustainability of wild populations, preserving the biodiversity of the wild salmonids of the Foyle system is therefore a primary objective of the Loughs Agency.

## 14.0 POLLUTION MONITORING

The Loughs Agency has a statutory obligation to monitor the pollution of watercourses. In conjunction with the Environment and Heritage Service and Louth County Council all reported pollution incidents are investigated.

## 15.0 FISHERY OFFICERS CARLINGFORD AREA REPORT 2008

In 2008 the number of agricultural related problems appeared to be less in 2008 than in previous years. This is most likely due to potential ramifications on single farm payments which can be affected by poor farming practice impacting upon the environment. Genetic sampling was conducted on the Atlantic salmon population of the Whitewater catchment as part of a wider project designed to provide a baseline of Atlantic salmon genetic structure throughout Northern Ireland.

As outlined above the Loughs Agency co-funded habitat improvement work on the Kilbroney River.

The Loughs Agency in conjunction with NIEA, Mourne Heritage Trust and Tara Seal Research undertook a comprehensive seal counting programme which will continue in 2009. This project will provide important information for assessing seal/fishery interactions within the Carlingford area.

A survey of Derryleckagh lake was conducted in 2008 and it is hoped that more lakes in the area can be surveyed in the future to provide important information on fish populations, distribution and age structures.

## 16.0 WHITEWATER RIVER SALMONID POPULATION SURVEY 2008

In September 2008 a quantitative electrofishing survey was conducted on the Whitewater River downstream of Ballyardle Bridge. This area is known to be stocked with fry and ova from the Whitewater hatchery. It is hoped that in future years that both stocked and non stocked sites will be able to be surveyed to provide an insight into the impact of stocking and the impact of natural recruitment on salmonid survival and growth. Basic biological data including densities, lengths and weights have been recorded and will provide a basis for comparison over coming years.

All rivers have a natural carrying capacity which means that above a certain threshold natural limiting factors such as food availability and size of feeding territories impact on the rivers ability to sustain more fish numbers. This is an extremely important factor when considering stocking locations and densities. Carrying capacities will vary depending on the type and location of the river. It is vital that any stocking if deemed necessary should be conducted in areas where no natural recruitment has taken place as this will increase the competition with naturally spawned fish which may be genetically “fitter”.

It was also noted in 2008 that there appears to be a bottleneck in natural production within the Whitewater River that is potentially caused by the disconnection of in-channel habitat caused by the significant numbers of weirs. The impact of these weirs could be two fold, firstly they can impact on migration as some are quite high with insufficient depth in the pool immediately downstream and secondly weirs act as a barrier to the recruitment of substrate from upstream. Rivers in their natural state will erode and accrete providing clean silt free gravel and nursery stones which will ultimately provide ideal habitat for salmonids and other native fish species.




Fig 16 Salmonid densities at Ballyardle Bridge 2008




Fig 16.1 Salmon fry and parr length weight relationships at Ballyardle bridge 2008 N = 89




Fig 16.2 Length Frequency distribution of Whitewater salmon fry and parr in 2008 N = 89



## 17.0 ACTIONS FOR 2009

In order to fully utilise the extensive data resources collected and held by the Loughs Agency on the fish populations and habitats of the Carlingford area it is necessary to focus attention on specific management objectives.

The Loughs Agency has stated in its corporate plan 2008-2010 that it will conserve, protect, manage and improve the fisheries of the Foyle and Carlingford areas. By way of fulfilling these objectives a targeted series of actions utilising data collected over recent years will be implemented. Fishery owners and local angling clubs will continue to be consulted regarding any proposed works and stakeholder input sought.

### 17.1 Foyle and Carlingford Areas Ongoing Actions for 2009

**Good water quality** is essential for the conservation of productive aquatic ecosystems. Fish populations rely on unpolluted water for survival and feeding. The Loughs Agency is committed to ensuring deleterious matter does not enter any watercourse. Routine monitoring is conducted throughout the Foyle and Carlingford areas. Proactive pollution visits and water quality monitoring will continue in 2009.

**Water quantity** is becoming an increasingly important issue from a fisheries management perspective with continuing demand from a variety of sources including industry, hydro power generation and abstraction for meeting the ever growing needs of industry and the wider population. The Loughs Agency are aware of the conflicting needs of aquatic environments and water resource users and comment on development issues which may have an impact on the important aquatic resources of the Foyle and Carlingford areas with reference to national and international obligations.

**In-channel and riparian habitat improvement** projects provide an important mechanism by which to improve and protect valuable fishery resources. Over recent years the Loughs Agency has developed a number of projects designed to improve the survival and production of robust populations of juvenile salmonid and other native fish species. These programmes will continue where funding is available. The Loughs Agency also encourages local stakeholder groups to source appropriate funding to develop collaborative habitat improvement projects. The Loughs Agency can provide advice and recommendations for in-channel and riparian improvements and are eager to facilitate the development of such programmes.

Work is continuing to assess and record all **Barriers to Migration** within the catchments of the Foyle and Carlingford areas and these will be incorporated into

the Loughs Agency Geographical Information System (GIS). Where finances are available the removal of artificial barriers will be investigated.

**Predation by cormorants and seals** of economically important fish species continues to be a contentious issue. The Loughs Agency will continue to promote the development of a management strategy incorporating economic, social and environmental factors.

**The Loughs Agency will continue to monitor the salmon and inland fishery resources of the Foyle and Carlingford areas**, utilising best practice methods including fish counters, juvenile population surveys and catch returns. The importance of the Atlantic salmon resource has been further highlighted by recent genetic studies which have identified the presence of genetically distinct populations of salmon between and within main river catchments. This information will be utilised when developing habitat improvement programmes to ensure the presence of a diverse resource capable of withstanding change.

**Invasive species** in both aquatic and riparian habitats have become an important issue in fisheries management and in wider environmental management. Invasive species have the potential to significantly alter ecosystems and their function. The Loughs Agency is contributing towards the development and implementation of invasive species codes of practice.

## 17.2 Carlingford Area Specific Actions for 2009

- Target all areas/individuals brought to Loughs Agency attention
- Implement habitat improvement schemes as dictated by business plan/corporate plan and the availability of resources
- Conduct annual fish population surveys and spawning surveys
- Conduct ongoing water quality monitoring and investigate areas highlighted as being of concern
- Develop potential habitat improvement projects including riparian buffer zone creation, fencing, native species planting and in-channel habitat improvements including spawning bed and nursery habitat improvement and identify resources
- Monitor forestry operations adjacent to watercourses or areas likely to impact on watercourses

- Assist with Water Framework Directive fish monitoring programme
- Monitor all sand and gravel extraction sites and onsite water management practices
- Ensure all fish passes, dams and mill races meet required standards
- Investigate habitat connectivity on the Whitewater River
- Instigate revised habitat survey on the Whitewater River
- Investigate fish passage issue on the Cassey Water
- Conduct genetic sampling of the Carlingford trout populations
- Continue to develop coarse fish survey programme where resources are available